

CIP GENERATION PROJECT
2013 COMMUNITY BENEFITS PROGRAM
REEF FISH MONITORING PROJECT
YEAR 6 RESULTS

Prepared For:

Environmental Department
Hawaiian Electric
P.O. Box 2750
Honolulu, Hawaii 96840-0001

Prepared By:

Richard Brock, Ph.D.
Environmental Assessment, LLC
1232 Lunalilo Home Road
Honolulu, Hawaii 96825

EA,LLC Report No. 2014-05

July 2014

EXECUTIVE SUMMARY

The development of an electrical generating facility at Campbell Industrial Park (CIP) Barbers Point was the impetus to initiate a quarterly environmental monitoring program to follow changes, if any, in coral reef fish communities in the Barbers Point - Kahe Point area. This document is the sixth annual report for this effort covering the period from December 2007 through December 2013 with a focus on the surveys completed in 2013. On a quarterly basis, this study monitors the status of coral reef fish communities at sixteen permanently marked sites offshore of Barbers Point on the southeast to Nanakuli Beach Park about 7.9 km to the northwest. These monitoring stations are all in waters from 5 to 12 m in depth and thus are subject to impact from high surf events.

Because of Hawaiian Electric's construction/operation of the generating station at Kahe Point as well as the developments at West Beach and Barbers Point Harbor, long-term marine environmental data covering the status of fish and coral communities are available commencing from the mid-1970's up to present. The most comprehensive of those efforts occurred with the Hawaiian Electric Environmental program in support of the Kahe Generating facility at Kahe Point. The Hawaiian Electric monitoring program documented changes that occurred to marine communities following three major storm events: the January 1980 event, Hurricane Iwa in November 1982 and Hurricane Iniki in September 1992 all of which severely impacted coral reef communities in the area. These studies demonstrated the impact of those storm events and not the operation of the Kahe facility as the major source of impact to marine communities of the Kahe area.

In the present study there were no statistically-significant changes in the mean number of fish species, mean number of individual fish censused or in the mean standing crop per transect among the twenty-four 2007-2013 survey periods, thus demonstrating stability in these communities. All species of fishes censused in the present study have been assigned to one of five feeding guilds (or trophic categories): herbivores (species feeding on algae), planktivores (species that feed on zooplankton up in the water column), omnivores (species that feed on both algae and small animals), coral feeders which are a specialized group feeding on coral tissue or mucous, and carnivores which are species feeding on smaller fishes and invertebrates living on the coral reef. Of the 146 species of fishes encountered in the twenty-four 2007-2013 surveys, twenty-three species are herbivores, fifteen are planktivores, seven are omnivores, eight are coral feeders and 93 are carnivores. Fifteen of the sixteen monitored locations are established on natural substratum where 87% of the fish standing crop is comprised of herbivores and carnivores. However, at one station established on the Kahe Generating Station warm-water discharge (Station 16), herbivores are largely replaced by planktivores but carnivores remain important as elsewhere. The reasons for this shift in dominance is due to the thermally-elevated discharge creating a unidirectional current that discharges particulate materials. The steel and armor rock covering the discharge pipe also provides a high degree of shelter space at this station.

This study was undertaken to follow changes in coral reef fish communities as part of the environmental monitoring program related to the development of the CIP electrical generation facility. The data collected in the first year represent the preconstruction baseline (December 2007 - December 2008), while data collected in the second year represents the construction phase (January - September 2009) and the data collected in the third through sixth years (2010-2013) represent the operational phase of the plant. The sixteen stations geographically fall into four groups along the 7.9 km of coastline; on the southeast are four stations offshore of the generation plant at Campbell Industrial Park (Station nos. 1-4), three stations seaward of Ko'Oolina Resort (nos. 5-7), five stations fronting the Kahe generation facility (nos. 8-12), three stations north of Kahe Point (13-15) and the Kahe Station discharge pipe (no.

16). Statistical analysis of the fish community parameters measured in this study (i.e., number of species, number of individuals and standing crop) on natural substratum found that the diversity of fish species, the number of individual fish as well as the standing crop to be significantly greatest at the three Ko'Olina stations over those in the other three groups over the 2007-2013 survey period. These three measures were least (or nearly so) at the four stations offshore of Campbell Industrial Park and these differences are attributed to better benthic community development offshore of Ko'Olina than elsewhere. The data from stations offshore of Kahe and from those to the north were between the Ko'Olina and Campbell Industrial Park means. The above analysis excluded data from Station 16 (the Kahe discharge pipe) because it is a man-made structure and not comprised of natural substratum as present at all other stations. However, to better understand the differences among the sixteen stations, the three fish community measures (mean number of species per transect, mean number of individuals per transect and mean estimated biomass per transect) were statistically examined comparing all stations. Two findings emerge: (1) the Kahe discharge pipe station had a non-significant greater number of species present and a clearly-separable significantly greater mean number of fish individuals and standing crop over all other stations and (2) the means for all parameters from all other stations were not statistically separable except for one station (number 11) where the number of fish species was significantly less than all other stations which is due to the short transect length sampling this site (i.e., 10.5 m versus 50 m) at all other sites. Thus the development in the fish communities at the fifteen stations situated on natural substratum monitored in this study pales relative to that found at the man-made Kahe discharge pipe.

Seven of the permanently marked monitoring stations in this study have been used in earlier Hawaiian Electric studies and the methods used herein are similar, allowing comparative analysis of the data. Comparing earlier fish community data (1976-1984) to present (2007-2013) data finds that there are no statistically significant differences in the annual mean number of fish species or annual mean number of individual fish censused per transect despite the imposition of three major storm events in 1980, 1982 and 1992 suggesting that the fish communities have to some extent recovered from these disturbances. These documented storm events impacted marine communities offshore of the Barbers Point and Kahe Point areas. These impacts were probably greatest on the coral communities which are the source of much of the natural local topographical relief creating shelter for fishes. If disturbance to the coral community occurs frequently and corals are known to be slow-growing, they are unable to contribute much to the local topography upon which many fish species depend thus keeping the fish community at an earlier point in community succession. The early studies demonstrated the large impact that these storms had at the time on corals as well as the movement of sand away from the Kahe area leaving much near-barren limestone that is present today and is scoured by small wave events keeping benthic community development to a minimum. This has resulted in a relatively poor development of the fish communities at many of the Kahe sample sites which continues to today. Where topographical relief and benthic communities are well-developed, the fish communities are likewise better developed. Given the long-term data set spanning 37 years and the apparent lack of strong significant changes occurring to fish communities with the three early storm events which is probably related to some level of recovery, suggests that the variation seen in the measures of the fish community used in this study will continue to fluctuate at a similar magnitude in future monitoring events as this program moves forward. Furthermore, the analysis of the 2007-2013 data suggest that benthic community development/topographic complexity creating shelter for fishes remain the overriding factors determining the degree of development in fish communities at the stations monitored in this study. Since these factors were heavily impacted by the early storm events many years ago (as documented by Hawaiian Electric), the present findings will probably continue much the same in future years of this study.

INTRODUCTION

1. Purpose

Hawaiian Electric has constructed a new generating station on vacant portions of its existing Barbers Point Tank Farm in Campbell Industrial Park (CIP) on the island of O'ahu. This generating facility was constructed in light of the fact that there is an urgent need for new generating capacity on the island. Initially, the generating station would consist of a single 110 megawatt (MW) Siemens-Westinghouse combustion turbine (CT) and two single 2 MW capacity black-start diesel engine generators. The system was designed to be fueled primarily by biofuels which assists in fulfilling the State's goals of energy security and sustainability. However, alternative fuels (e.g., diesel, naphtha, etc.) may be used if biofuels are unavailable. The facility is designed to accept a second generating unit and could be constructed if and when it is needed to meet system requirements. It was expected that the generation system could be used to help meet peak load periods on the island's system which normally occur between 5:00 pm and 9:00 pm on weekdays.

The single CT generation unit utilizes approximately 600 gallons per minute (gpm) of water which is used for water injection into the CT for air pollution control, equipment cooling, plant washdown, landscape irrigation and domestic use by operating personnel. Disposal of used water is via injection wells on the facility site. Thus, unlike the nearby Kahe Generating Station where seawater is used for cooling in the plant and discharged back into the marine environment, the CIP plant does not discharge cooling water into the nearby ocean thus precluding or significantly reducing potential for environmental impacts to occur in the marine environment.

As part of the environmental monitoring program for the CIP Generating Station, it was suggested that a coral reef fish monitoring program be put in place to track the changes, if any, that may occur with fish populations offshore of the plant at Barbers Point. The data presented were collected in 2013. Data were initially collected in 2008 representing the preconstruction baseline, in 2009 representing the "construction period" of the generating facility and in 2010 representing the commencement of the operational phase of the plant and continuing through 2013. The 2008 information was presented in Brock (2009), the "during construction" information was given in Brock (2010) and data collected since the commencement of plant operations in 2010 is given in Brock (2011), for 2011 in Brock (2012), 2012 in Brock (2013) and the continuing operational phase data for 2013 are presented herein.

Since Hawaiian Electric had this type of monitoring plan in place offshore of its Kahe power plant in the 1970's and 1980's, the present study has included a reassessment of some of those locations which provide information on the changes that have occurred to fish communities in the Barbers Point - Kahe Point area over the last 30 years. This study addresses the question, "What are the changes in the coral reef fish community structure that occur through time in the Barbers Point - Kahe Point area?" Community structure is defined as the diversity of species,

their abundance and biomass as well as their place in the food web of the coral reef. This document addresses this question and represents the sixth annual report since monitoring began for the CIP Generating Station and the fourth annual report since it began operating. This report includes a comparative assessment to baseline and during plant construction periods.

2. Natural Events and Impacts to Hawai'i's Coral Reefs

It is a common belief that coral reefs and their fish communities exist in stable environments which have resulted in the high diversity of species that is often seen in these systems. More recent data has shown that the environment in which coral reefs exist is dynamic, i.e., undergoing constant change, thus the organisms are subjected to a variety of stresses, resulting in shifts in community structure and abundance of species (Grassle 1973, Connell 1978, Dollar and Tribble 1993). Indeed, the concept that “intermediate levels of disturbance” may result in higher diversity has been demonstrated in a number of studies of coral communities (Connell 1978, Dollar 1982, Grigg 1983). Benign environments result in final successional stages of coral community development with low species diversity where one or just a few species dominate. This decrease in species diversity is found also with the coral-associated fish communities. Stability in coral species populations has been recently viewed as ever-changing in time and space, where species diverge by genetic drift due to isolation or converge by hybridization, producing constant change which has been described as reticulate evolution (Veron 1995).

Stochastic (i.e., random) processes create a nonequilibrium situation in coral reef communities. A major causal mechanism of stochastic events is the occurrence of occasional storms, which have been shown to be the single most important factor influencing the structure, diversity, and abundance of coral communities in Hawai'i (Dollar 1982, Grigg 1983, Dollar and Tribble 1993). Coral reefs have been described as “temporally varying mosaics” (Bak and Luckhurst 1980) in which the coral community undergoes a continual cycle of disturbance or removal and recovery or renewal. The effects of severe disturbance that drive this cycle have been documented for specific reef areas. The removal or destructive phase due to large storm events has been recorded in the Caribbean (Ball *et al.* 1967, Perkins and Enos 1968, Stoddart 1969, 1974, Woodley *et al.* 1981) and in the Pacific (Blumenstock *et al.* 1961, Cooper 1966, Dollar 1982, Dollar and Tribble 1993, Done *et al.* 1991, Harmelin-Vivien and Laboute 1986, Maragos *et al.* 1973, Ogg and Koslow 1978).

Following the impact of large storm events that disrupt the coral and fish communities is a period of regrowth. This period has received less study because the recovery of most coral communities is a slow process and because having pre-storm study sites where post-storm sampling can be done is rare (Dollar and Tribble 1993). Corals are relatively slow-growing and long-lived, thus the successional processes on most reefs take place on a scale of years to decades (Grigg and Maragos 1974).

In exposed locations in Hawai'i, storm waves keep coral communities at an early point in

succession (Dollar 1982, Grigg 1983, Dollar and Tribble 1993). Under such situations, coral colonies never attain any significant size and growth forms are usually prostrate, thus reducing their exposure to wave energy. Since much of the development in the associated fish community is related to the topographical complexity of the substratum (Risk 1972) and much of this complexity is directly due to the growth of corals, fish community development is usually reduced where coral communities are poorly developed and shelter space is lacking. Besides topographical complexity providing shelter habitat for fishes, the highly variable shelter created by coral communities serves a wide range of invertebrate and algal communities which may be forage for many fish species. Thus the development of coral reef fish communities is often directly linked to the degree of development of coral communities and factors that negatively affect the coral community frequently will have a similar negative impact to the fish community.

In general, many corals in Hawai'i have relatively slow growth rates, and many species produce annual growth bands much like the large conifers of temperate forests (Knutson *et al.* 1972). The large hemispherical colonies of *Porites lobata* do this, accreting about a centimeter per year in radial diameter. In Hawai'i, *P. lobata* colonies may attain diameters in excess of 4 m, thus large colonies may be more than 150 years in age. Under these circumstances, significant storm events do not have to occur with much frequency to have a strong influence on the successional state and development of coral communities where this species occurs.

Since 1980, three major storm events have created large surf that have caused impacts to Hawai'i's reefs over levels that normally occur. The January 1980 storm brought waves which attained heights of at least 6 m, from a south-southwest direction to the islands (Dollar 1982) thus impacting the Barbers Point - Kahe Point region. The next major storm event was Hurricane Iwa, which struck the islands in November 1982. Again, storm waves which attained estimated heights of 9 m, impacted the south and west shores of all islands (Coles and Fukuda 1984). The most recent major storm event was Hurricane Iniki, which passed over Kauai on 11 September 1992 with sustained winds up to 144 miles per hour (mph). It also created large surf that again impacted the south and west shores of O'ahu with storm generated surf arriving from a south-southeast (SSE direction). On the south shore of O'ahu, wave heights were estimated to reach 8 m (Brock, personal observations).

3. Hawaiian Electric's Environmental Monitoring Program: A Synopsis of Impacts Associated with the Construction and Operation of the Kahe Generating Station (1970's-1980's)

As part of the National Pollutant Discharge Elimination System (NPDES) permit conditions allowing the discharge of thermally-elevated cooling water into the marine environment at Kahe Point, Hawaiian Electric was required to monitor the status of the coral, algae and fish communities in the offshore waters fronting and in the vicinity of the plant. The findings from these early monitoring efforts provide an excellent overview of the environmental changes that occurred in the Kahe Point area prior to the three storm events in 1980, 1982 and 1992 and after the January 1980 and November 1982 events. Studies on coral coverage showed a significant

decrease of 7% from 1973 to 1975 and an additional 13% from 1975 to 1977. These decreases were correlated with proximity to the Kahe plant discharge but the analyses did not determine whether the disturbance associated with outfall construction or plant operation were definitive factors producing the mortality. In contrast to the increased mortality, settlement and growth of coral recruits increased with proximity to the outfall after the plant began operating, which suggests that outfall construction rather than plant operation was the major factor in producing the mortality. Fish populations throughout the study area showed no changes except on the marginal reefs to the northeast of the outfall where both the numbers of species and individuals censused decreased following the commencement of outfall operations. However, the number of intertidal species on the rocky shoreline increased in the areas of thermal impingement (Coles *et al.* 1985a).

In 1978 the analysis of all reef fish population data collected since the beginning of the offshore outfall operation in December 1976 indicated that fish populations were being displaced from the immediate vicinity of the outfall (Coles 1979). The impact caused by the January 1980 “Kona” storm that generated extreme surf on the south and western shores of the islands, however, was much greater than the changes observed from 1976 to 1978. The Kahe study area was heavily impacted by waves at that time. Subsequent survey work found that the 1980 storm was responsible for reductions in coral coverage, fish populations and the redistribution of beach sand that were all much greater than the subtle changes which had occurred in these parameters over the previous seven years (Coles *et al.* 1981).

During 1981, the generating capacity of the Kahe Station was increased by the addition of Unit 6 to a total of 651 MW which increased the cooling water flow to 846 million gallons per day (mgd), a 33% increase above the flow rate for Units 1 to 5. With this change came a reduction in the surface plume area to about one-half while the area of benthic thermal impingement nearly doubled, but was restricted primarily to offshore sand areas. A result of these changes was a moderation in coral coverage declines seen previously but coral reef fish populations continued to decline probably in response to the decrease in reef habitat produced by the 1980 storm (Coles *et al.* 1982).

In November 1982, Hurricane Iwa struck the Hawaiian Islands with most of the damage occurring on Kauai. On O'ahu, damage was greatest along the northwest coastline which included the Kahe Point area. Waves and winds were substantially greater than seen in the January 1980 event with waves heights estimated at 30 feet (Noda 1983). As described in Coles *et al.* (1985a, page 16):

“Surprisingly, coral communities in shallow water areas appeared relatively undisturbed by hurricane wave turbulence. However, reefs further offshore at depths of 20 feet or more appeared to have been substantially destroyed by the force of breaking waves. Measurements of reef coral coverage and fish populations just prior to the hurricane had indicated stable populations compared to the previous year, indicating that damage had resulted from the catastrophic forces released by the hurricane. A further observation of interest was that sand

along the reef front had been swept away by the hurricane's waves, exposing reef pavement and rubble that had been buried by up to five feet of sand.

The 1983 monitoring investigations verified the preliminary conclusions that had been determined shortly after Hurricane Iwa occurred. Quantitative estimates indicated substantial reductions in coral, algal and fish communities corresponding to locations where hurricane wave forces had been greatest. Due to removal of sand from shallow areas and the extreme cutting back of beaches that had occurred during the hurricane, sand entrainment through the Kahe Station was substantially less in 1983 than during previous years. A study of coral recolonization in the area indicated a positive influence of the Kahe outfall in the re-establishing of reef corals on denuded reef surfaces."

Coles and Fukuda (1984) noted the net significant decrease in coverage of 18.7% between 1979-1980 due to the January 1980 storm as measured at the Kahe permanent monitoring stations. Hurricane Iwa contributed a further significant decline of coral offshore of the Kahe facility; in 1982-83 the net change in coral coverage decreased 5.4%. The greater decline in coverage with the 1980 storm relative to Hurricane Iwa was probably related to two facts: (1) since the wave energy of the January 1980 event was less than the 1982 hurricane, the impact of that energy was probably released at shallower depths where coral coverage had been high and (2) Hurricane Iwa occurred just two years after the January 1980 storm event leaving little time for significant coral recovery to occur.

4. The Impact of Hurricane Iniki

As noted above, Hurricane Iniki struck the Hawaiian Islands in September 1992 with high waves impacting the south and west shores of all islands. Fifty-four days after Hurricane Iniki, a qualitative survey was carried out to determine the extent of damage to coral communities in the vicinity of the Kahe Generating Station (Brock 1992a). Fourteen of the more than 38 permanently marked monitoring stations were visited. With respect to coral damage, two general findings emerged: (1) that damage due to storm waves to corals was minimal and was primarily restricted to the cauliflower coral *Pocillopora meandrina*. The reasons for this restricted damage was related to the branching nature of this species as well as the fact that this coral frequently colonizes the tops of high points on hard bottom (i.e., limestone ridges and boulders). In these locations, cauliflower corals have relatively greater exposure to wave energy impinging on the bottom than would coral colonies situated down in depressions. The second finding was that the greatest damage to corals occurred at those stations situated in areas with greatest exposure to wave forces impinging from the SSE direction which was consistent with the direction of Hurricane Iniki's storm waves. Finally, the field survey noted that a considerable amount of sand was removed by the storm at some stations with a net result of a greater amount of hard substratum previously covered by sand was now exposed and available for benthic recruitment. Only one station examined in the study showed evidence of net deposition of loose materials (i.e., coral rubble and broken live pieces) while at all other stations, sand, broken live corals and rubble were not present and assumed to have been advected to deeper water seaward and outside

of the study area (Brock 1992a). These findings were similar to those noted in Mamala Bay, southeast of Kahe study area (Brock 1996).

As noted above, Hawaiian Electric carried out environmental surveys following the January 1980 storm and Hurricane Iwa in 1982. Several key observations emerge in comparing the findings following the 1980 storm to those from the post-Hurricane Iniki study: the January 1980 event had a much greater impact to the Kahe coral communities relative to Hurricane Iniki; (2) it caused considerable deposition of sand at many stations which in some cases caused burial of corals; and (3) it was responsible for significant abrasion of many corals which was not obvious following Hurricane Iniki. The finger coral, *Porites compressa*, was present at many of monitoring stations in 1980 and by the time of the post-Iniki survey, this species contributed little to the coverage estimates at sampled stations. Because of its relatively delicate skeletal structure, *P. compressa* is prone to damage by storm surge (Dollar 1982) and the storms since 1980 have probably contributed to the decline of this species at many Kahe Point locations (Brock 1992a).

The energy from the high amplitude, short period waves generated by all three storm events (January 1980, November 1982 and September 1992) was dissipated in deeper water thus coral communities in these deeper areas were potentially exposed greater impact (see Dollar 1982, Walsh 1983). As noted by Coles and Fukuda (1984), fully 90 percent of the coral coverage offshore of the Kahe generating facility was at depths of 10 m or more prior to the January storm. These deeper water coral communities apparently received much of the damage in 1980 and again in 1982 with much of that damage occurring to the finger coral, *Porites compressa*. Brock (1992b) examined marine communities southeast of the Barbers Point Deep Draft Harbor two weeks after Hurricane Iniki and found considerable damage to corals below 13 m and the damage was greatest in areas exposed to a SSE swell. Coral communities inshore of this or those protected from a direct SSE swell direction, appeared to have suffered little impact. Brock's observations included the disappearance of a large amount of loose coral rubble in the 12 to 22 m depth range where rubble that had accumulated intermittently along the base of a submarine cliff. Individual estimated volumes were in excess of 2,000 cubic meters (m³) over linear distances of 30-50 m and this material was not found within diving depths (here from shore to 30 m).

As noted by Brock (1992a, page 5):

"The two storms preceding Hurricane Iniki produced opposite impacts subtidally with respect to the movement of sand offshore of the Kahe facility. The January 1980 storm resulted in the deposition of sand over many reef areas, thus burying or scouring benthic communities. In contrast, Hurricane Iwa resulted in 3 to 5 feet of sand being removed along the seaward edge of the reef exposing coral reef framework that had been formerly covered. Coles and Fukuda (1983) noted '...sand which had been deposited by the Kahe outfall and swept on to the reefs by previous storms was completely removed from along the entire reef front. The substratum available in the area is now similar to the conditions when marine monitoring began in 1973...'. It appears that Hurricane Iniki also removed sand from the area seaward of the forereef but to a much lesser extent than in the November 1982 event (i.e., up to 0.75 m in 1992 versus up to 1.5 m

in 1982); perhaps the sand had not returned before the 11 September 1992 storm.”

The three strong storms commencing in January 1980 and ending 12 years later with Hurricane Iniki documented change to the bottom communities in the Barbers Point - Kahe Point area. These changes also created a negative impact to the resident fish communities which has been documented elsewhere in Hawai'i (Walsh 1983). The findings from these past studies, therefore, indicate that knowledge of the past environmental history can lead to a better understanding of the biological resources present in the area today. This environmental history provides the basis for the present study.

METHODS

The fish communities at sixteen permanently marked sites are monitored on a quarterly schedule. These sixteen sites are located in the Barbers Point to Nanakuli area on the west coast of O'ahu (see below). The monitoring of fish communities is carried out using a visual census method. The sampling protocol occurs in the following sequence: on arrival at a given station, the individual conducting the visual fish census enters the water and carries out the visual census over a 50 m long by 4 m wide corridor run parallel to shore. (Only Station 16, which is located on the Kahe facility's discharge pipe, runs perpendicular to shore and station 11 is only 10.5 m in length). All fishes within this area to the water's surface are counted. Data collected include the species, numbers of individuals and an estimate of the length of each individual fish counted. The length data are later converted to standing crop estimates using linear regression techniques. The diver equipped with SCUBA, transect line, slate and pencil enters the water, counts and notes all fishes in the prescribed area (method modified from Brock 1954). The 50 m transect line is paid out as the census progresses, thereby avoiding any previous underwater activity in the area which could frighten wary fishes. The length data are used in making estimates of biomass for each species present coupling the length data with species-specific regression coefficients (Ricker, 1975, Brock and Norris 1989).

Fish abundance and diversity are often related to small-scale topographical relief over short linear distances. A long transect may bisect a number of topographical features (e.g., cross coral mounds, sand flats and algal beds), thus sampling more than one community and obscuring distinctive features of individual communities. To alleviate this problem, a relatively short transect (50 m in length) has proven adequate in sampling many Hawaiian benthic communities. In addition, the transect length used by Coles *et al.* (1985a) was also 50 m thus making the present counts collected under this program comparable to the earlier data collected by Hawaiian Electric. However as noted above, Station 11 which was originally established by Hawaiian Electric in the 1970's is only 10.5 m in length.

Besides frightening wary fishes, other problems with the visual census technique include the underestimation of cryptic species such as moray eels (family Muraenidae) and nocturnal species, e.g., squirrelfishes (family Holocentridae), aweoweos or bigeyes (family Priacanthidae), etc. This problem is compounded in areas of high relief and coral coverage affording numerous

shelter sites. Species lists and abundance estimates are more accurate for areas of low relief, although some fishes with cryptic habits or protective coloration (e.g., the nohus, family Scorpaenidae; the flatfishes, family Bothidae) might still be missed. Obviously, the effectiveness of the visual census technique is reduced in turbid water and species of fishes which move quickly and/or are very numerous may be difficult to count and to estimate individual sizes. Additionally, bias related to the experience of the diver conducting counts should be considered in making any comparison between surveys. In spite of these drawbacks, the visual census technique probably provides the most accurate nondestructive method available for the assessment of diurnally-active fishes (Brock 1982).

In the analysis of the data, all fishes encountered were classified as to their primary foraging behavior as a means to better understand the trophic relationships in the fish communities. These functional groups are carnivores which includes all fishes feeding on other coral reef animals (fish and invertebrates) greater than zooplankton in size, planktivores which are species that feed primarily on zooplankton and detritus in the watercolumn, herbivores which are species feeding primarily on algae, omnivores which are usually small species that feed on a combination of algae and benthic animals and the coral feeders which are a specialized group of fishes that feed on coral polyps and mucous. The determination of which species were in each feeding guild utilized the findings of Hiatt and Strasburg (1960), Hobson (1974), Brock *et al.* (1979) and Randall (2007). Primarily nonparametric statistical procedures are used thus avoiding the requirements for normality in the data, etc. that are necessary in parametric statistical analyses.

RESULTS AND DISCUSSION

1. Station Locations

To assess the status of coral reef fish communities in the Kahe-Barbers Point area, sixteen permanently marked stations were established. These stations are spread along 7.9 km (4.9 miles) of coastline fronting the CIP Generating Station at Barbers Point on the southeast to the south boundary of the Nanakuli Beach Park on the northwest and their approximate positions are shown in Figure 1 and more precise locations (latitude/longitude) are given in Table 1. Eight stations were established in 2008, prior to the pre-construction monitoring event and the rest are stations established for the Hawaiian Electric environmental monitoring program in the 1970's. Four stations are located offshore of Campbell Industrial Park at Barbers Point in waters from 7 to about 10 m in depth. These stations (Station nos. 1 - 4, Table 1) monitor the status of fish communities in closest proximity to the CIP Generation site and are located to the southeast of the Barbers Point Harbor entrance channel. Two stations are located northwest of the Barbers Point Harbor entrance channel fronting the Ko'Olina Resort and Paradise Cove area (Station nos. 5 and 6, Table 1). Again the water depths at these two stations is from 7 to 9 m. Coles *et al.* (1985) monitored fish community structure at seven stations fronting and adjacent to the Kahe Generating Station. These seven stations are also monitored in the present study (here numbered as Station nos. 7 through 13 in Table 1) to obtain information on the status of these fish communities today and to compare the fish community structure today to what was present at

these same locations more than 25 years ago. These stations are in water ranging from 5 m to 12 m in depth.

The previous Hawaiian Electric environmental monitoring program also monitored a control station offshore of Nanakuli (Coles *et al.* 1985a) which has also been added to the stations monitored under the present program (here Station 14, Table 1). A second control station (Station 15, Table 1) approximately 70 m north of Station 14 has been established for the present monitoring program. Finally Station 16 was established on the Kahe discharge pipe directly offshore of the generating facility in water from 5 to 7 m in depth.

As noted above, the locations of all stations are shown in Figure 1. The “start point” for each station is marked using 90 cm long nylon cable ties and small subsurface floats that are tied to the substratum in proximity to the start point for each transect. Because of high public use by dive tour operators and individuals SCUBA diving from shore fronting the Kahe Generating Station, Stations 7 - 12 as well as Station 16 have not been marked but rely on prominent natural points on the local substratum. Past experience in permanently marking biological monitoring stations in “high use” areas results in divers removing materials of anthropogenic origin thus destroying and negating this method for relocation of stations. Low-cost modern global positioning systems (GPS) can put the diver/monitor within a few feet of any known point. The GPS waypoints for each of the 16 stations sampled in this study are given in Table 1.

2. The 2007-2008 (Preconstruction) Data

During the preconstruction period, fish transect data were collected on five occasions commencing on 27 December 2007. In 2008, transect work was carried out on 4 April, 30 May, 19 August and on 25 November. As noted above, sixteen stations were routinely sampled in this study and these early data are presented in Brock (2009). In the first survey, twelve of the sixteen stations were sampled; missing were Stations 4 (East 4), 5 and 6 (Ko'Olina 1 and 2) and 16 (Hawaiian Electric discharge pipe). The second survey carried out on 4 April only missed one site, Station 16 (the Hawaiian Electric discharge pipe) and by the third survey on 30 May 2008 all sixteen sites were sampled. The Hawaiian Electric thermally-elevated discharge (Station 16) was added as a monitoring station because of the well-developed fish community present at that location. Because station 16 is unusual with a highly developed community on a man-made structure, it is treated separately in many of the analyses below. In total, 122 species of fishes were censused in these first five surveys and these are given in Brock (2009).

3. The 2009 During Construction Data

In 2009 field surveys were conducted on 19 March, 11 May and 21 July. When the fourth quarter 2009 period commenced, weather deteriorated with a series of fronts that started in October 2009 and carried though unabated April 2010. Locally, these weather fronts brought surf as did weather fronts occurring elsewhere in the Pacific which affected the south, west, northwest and north coastline of O'ahu. Surf from these directions impinge on some or all of the

sample sites precluding field sampling during these periods. The result was that the fourth quarter 2009 field survey was not completed. Thus the analysis below includes data from the first three quarters of 2009 which represent the during construction period for the new generation facility at Campbell Industrial Park.

4. The 2010, 2011, 2012 and 2013 Operational Phase Data

In 2010 field surveys were carried out on 29 March, 14 May, 12 August and 29 October representing the first year of operations of the new generating facility at Campbell Industrial Park. Although the data collection phases have been split into “preconstruction”, “during construction” and operational periods, it should be noted that the CIP Generating Station is situated well inland of the ocean and its operation has no direct input to the sea. The 2011 surveys were carried out on 25 February, 16 June, 29 July and 23 November, the 2012 surveys were completed on 2 May, 23 May, 23 July and 2 November 2012 and the 2013 surveys were conducted on 3 May, 14 June, 20 September and 18 December 2013. Again, inclement weather (surf) precluded the timely collection of data in the first quarter for both 2012 and 2013 thus once the weather became favorable in late April 2012 and in May 2013, the first quarter field sampling was carried out on 2 May 2012 followed by the second quarter field work being done on 23 May 2012. Similarly, the first quarter 2013 field sampling was completed on 3 May and the second quarter work done on 14 June. The 2013 data are presented below along with a comparative analysis of all data collected to date.

The complete data set from the four 2013 surveys is given in Appendix 1 and this information is summarized in Table 2 along with the earlier (2007-2012) information. Drawing from some of these data and excluding Station 16, we may ask the question, “Are there any statistically significant differences among the mean number of fish species seen per transect, the mean number of individual fish censused per transect or the mean estimated standing crop in grams per square meter (g/m^2) among the twenty-four 2007-2013 sample periods?” To address this question two nonparametric tests were used: the Kruskal-Wallis analysis of variance (ANOVA) and the Student-Newman-Kuels (SNK) Test. The Kruskal-Wallis ANOVA is able to demonstrate statistically significant differences among parameter means (by date) but cannot show where those differences are. The SNK Test is used to group related sample means and separate those means that are significantly different from one another. The results of these analyses are given in Table 3. Referring to Table 3, the Kruskal-Wallis ANOVA noted no statistically significant differences exist among the means for each of the twenty-four sample dates for the number of fish species per transect, the number of individual fish censused per transect or for the mean estimated standing crop in g/m^2 per transect. These results point out that when considering grand means for the number of species, number of individuals or biomass (in g/m^2) per transect on each of the twenty-four sample dates, there are no significant differences. Thus at this level of analysis (i.e., grand means), there is no statistical separation among the dates which suggests a level of stability in the fish communities at these sample sites.

Station 16 established on the terminus of the Kahe Generating Station discharge pipe is

discussed separately because it is a man-made structure deployed in an area of sand bottom and has over 846 mgd of thermally-heated seawater discharging at its terminus. The topographical relief afforded by the steel and basalt rock substratum as well as coverage by corals is considerably more attractive to many fishes than the nearby surrounding natural reefs and the discharge of thermally-elevated water serves to attract many fishes. These features result in an enhancement of the local fish community making the structure of the fish community very different than that of any other of the fifteen natural reef sites sampled in this study. Thus as noted above, the results of fish censuses undertaken at Station 16 are discussed separately in most analyses.

The fishes censused in the twenty-four recent December 2007 - December 2013 surveys were assigned to one of five trophic categories or feeding guilds. As noted above, these groups are herbivores (species that feed on algae), planktivores or species that feed up in the water column on zooplankton, omnivores that feed both on plant material as well as small animals, coral feeders which are a specialized group feeding on coral tissue and mucous, and the carnivores which are species feeding on fishes and invertebrates found on coral reefs. In the five surveys carried out during the preconstruction (2007-2008) period there were 113 species of fishes encountered at the sixteen sample sites. The three during plant construction surveys completed in 2009 found 107 species of fishes at these sixteen sample sites. In the four operational phase 2010 surveys, there were 109 species of fishes recorded at the sixteen sites, in 2011, 106 species, in 2012, 100 species and in the four 2013 surveys there were 107 of fishes were sighted at the sixteen sites. In total among the twenty-four surveys, 146 species of fishes have been recorded among the sixteen survey sites. Fifty-four percent or 79 species encountered were in common among the twenty-four surveys carried out over the six-year period. These data suggest a reasonable level of stability in these fish communities.

Of the 146 species of fishes recorded over the twenty-four surveys, 64% (or 93 species) are carnivores, 16% (or 23 species) are herbivores, 10% (or 15 species) are planktivores, 5% (or 7 species) are omnivores and 5% or eight species are coral feeders. The assignment of fish species to the five trophic categories are given in Appendix 1 of this report as well as in Brock (2009, 2010, 2011, 2012 and 2013) for species encountered on each transect and earlier survey dates. Table 4 summarizes the feeding guild information by survey date providing the mean percent contribution by weight of each trophic category for stations in two groups; the first group includes stations 1 through 15 (natural substratum) and the second group considers only station 16 (the Kahe outfall station). Although the data in Table 4 are in summary form, two facts emerge, (1) that the majority (here 87%) of the weight of fishes censused at the first fifteen stations is comprised of herbivores and carnivores and (2) the importance of herbivores is largely replaced by planktivores at the Kahe outfall station (station 16, here 33.1%) but carnivores remain important at Station 16 (47.7%) as they are elsewhere. The large volume of thermally-elevated water (846 mgd) is probably serving both as a source of food (entrained particles that have passed through the plant) and a warm and strong unidirectional current that attracts and holds planktivorous species that naturally orient themselves into the current seeking food. In addition, the steel and armor rock superstructure that covers the Kahe facility's discharge pipe

along with high coral coverage provides habitat shelter and for some fish species a suitable substratum for spawning. A considerable part of the planktivore biomass at station 16 is comprised of two sergeant major or mamo species (*Abudefduf abdominalis* and the recently recognized *Abudefduf vaigiensis*) both of which not only feed in the discharge plume and environs, but also lay demersal eggs on the rocky substratum overlaying the discharge pipe. These two species along with the paletail unicornfish or kala lolo (*Naso brevirostris*) dominate the planktivore biomass at this site together comprising almost 41% of the total estimated standing crop present based on the 2008-2013 data.

4. Differences in Fish Community Structure in the Study Area

This study was undertaken to follow changes in coral reef fish communities as part of the environmental monitoring program related to the development of the CIP generation facility. Sixteen sites spread along 7.9 km of coastline are monitored (Figure 1); referring to Figure 1, these sites geographically fall into four groups: on the southeast are four stations offshore of Campbell Industrial Park and the generation plant (Station nos. 1-4 or East 1 through 4), three stations seaward of Ko'Olina Resort (Station nos. 5-7 or Ko'Olina 1 and 2 as well as Hawaiian Electric 1D), five stations fronting the Kahe generation facility (Station nos. 8-12 or Hawaiian Electric 5B, 7B, 7C, 7D, and 7E) and three stations to the north of Kahe Point (Station nos. 13-15 or Hawaiian Electric 10C, Nanakuli 1 and 2). Because Station 16 (the Kahe discharge pipe) is a man-made structure and not natural substratum like the other fifteen monitored sites, it is excluded from the present analysis.

The question, “Are there any statistically significant differences among the mean number of fish species per transect, the mean number of individual fish per transect or the mean estimated standing crop (in g/m²) per transect among the four above geographic groups of stations established on natural substratum and sampled in the 2007-13 period?” can be answered again using the Kruskal-Wallis ANOVA and the SNK Test. The results of these statistical procedures are given in Table 5. As noted previously, the Kruskal-Wallis ANOVA can discern whether means differ significantly but cannot separate those that are thus the SNK Test is used to demonstrate which means differ significantly from the others. In all cases the ANOVA noted significant differences exist among the four areas (i.e., CIP, Ko'Olina, Kahe and Nanakuli) for all three parameters (i.e., mean number of fish species per transect, mean number of individual fish per transect and mean standing crop of fishes per transect). The SNK Test demonstrated that the mean number of fish species, individuals and standing crops are significantly greater and statistically separable at the Ko'Olina group of stations over the three other station groups, all of which were related (Table 5). Coral community development (coverage) appears to be greater at the three Ko'Olina stations than found at any of the other transect sites and may be responsible for the greater diversity of species, numbers of individuals and standing crops present there.

Summarizing the results presented in Table 5, several (often non-significant) trends are apparent. First, the diversity of fish species, the numbers of fishes counted and their estimated biomass are greatest at the three Ko'Olina stations (Station nos. 5-7) and are least at the four

stations offshore of Campbell Industrial Park (Station nos. 1-4) for two of the three parameters. The development of benthic communities, including corals, is greater at Ko'Olina than it is offshore of Campbell Industrial Park where the topographical complexity which often serves as shelter for fishes is probably the least among the four station groups. Benthic community development, which includes the development of corals and topographical complexity, are probably less at the Kahe (Station nos. 8-12) and the North group (Station nos. 13-15) of stations relative to Ko'Olina but greater than found offshore of Campbell Industrial Park. Also, the mean number of fish species and individuals censused per transect was greater at the north group of stations (Station nos. 13-15) relative to the Kahe stations (Station nos. 8-12) and the CIP stations (Station nos. 1-4). However and as noted above, the mean estimated biomass of fishes per transect was significantly greater at the Ko'Olina stations over all other stations groups but the CIP group of stations has a non-significant greater standing crop than either the Nanakuli or Kahe station groups. The probable reason for the greater estimated standing crop at the CIP stations is the presence of a natural circular depression (about ten meters in diameter and one meter deep) having undercut ledges located about 6 meters shoreward of station 1. Because of the shelter created by this depression, a number of adult schooling surgeonfishes forage out across the substratum away from the depression and occasionally these fishes will pass through the transect during censusing resulting in higher biomass estimates.

The final statistical analysis of the 2007-2013 fish census data examines the mean number of fish species per transect, the mean number of individual fish per transect and the mean fish biomass per transect (in g/m²) examining each of the sixteen stations again using the Kruskal-Wallis ANOVA and the SNK Test. In this analysis, the question is “Are there any statistically significant differences between the mean number of fish species per transect, the mean number of individual fish per transect or the mean estimated standing crop among the 16 stations sampled in 2007-2013?” and the results are given in Table 6. Referring to Table 6, the Kruskal-Wallis ANOVA noted strong statistical differences among the means for all three parameters but the SNK Test found few clearly significant differences. These differences were: (1) the Kahe discharge pipe station has a greater (but insignificant) mean number of fish species, individuals (significantly greater) and standing crop (significantly greater) over all other stations and (2) the means from all of the other fifteen stations (located on natural substratum) are all related due to overlap in the SNK Test results except at station 11 where the mean number of fish species is significantly less than at all other stations (all of which are related). Station 11 also has the least mean number of individual fish (non-significant) and the second lowest mean standing crop (not statistically separable). These low numbers are probably related to the poorly developed coral community at station 11 resulting in little shelter present and most importantly, to the short transect length (10.5 m) at this station relative to all others which are 50 m in length. The obviously (non-significant) greater mean number of species, and significantly greater number individuals and standing crop at the Kahe discharge pipe is related to the presence of ample shelter, a unidirectional flow of thermally-elevated water and sufficient food resources present relative to all other stations which are located on natural substratum.

5. Fishery Resources

Appendix 1 in this report as well as in Brock (2009, 2010, 2011, 2012, 2013) provides lists of all fish species seen over the twenty-four 2007-2013 surveys. In these lists are both species that are sought-after by commercial, subsistence and recreational fishers as well as species that are usually not. In the usually sought-after group of species, most of the individual fishes encountered on the transects were juveniles but occasionally adult individual fishes were seen. Among the species seen include a number of small schools of the mackerel scad or opelu (*Decapterus macarellus*) especially around stations fronting the Kahe Generating facility in the December 2007 survey and scattered through the various stations and sample dates were seen adults of the moano kea (*Parupeneus cyclostomus*), omilu (*Caranx melampygus*), smaller individuals (papia) of the barred jack (*Carangoides ferdau*), lemon spot jack (*C. orthogrammus*), ulua aukea (*Caranx ignobilis*), pa'opa'o (*Gnathanodon speciosus*). Adults of other species seen include the lai (*Scomberoides lysan*), uku (*Aprion virescens*), wahanui (*Aphareus furca*), the introduced ta'aape (*Lutjanus kasmira*) especially at stations 13 and 16 and to'au (*Lutjanus fulvus*), weke (*Mulloidichthys flavolineatus*), weke'ula (*M. vanicolensis*), munu (*Parupeneus insularis*), moano (*P. multifasciatus*), malu (*P. pleurostigma*), kumu (*P. porphyreus*), nenu (*Kyphosus sandwicensis*), 'a'awa (*Bodianus bilunulatus*), kupoupou (*Cheilio inermis*), po'ou (*Oxycheilinus unifasciatus*), laenihi (*Iniistius umbrilatus*), the parrotfishes or uhus (*Scarus rubroviolaceus*, *S. psittacus*, *S. sordidus*, *S. perspicillatus*, *Calotomus carolinus*), the surgeonfishes including paku'iku'i (*Acanthurus achilles*), palani (*A. dussumieri*), maikoiko (*A. leucoparicus*), ma'i'i'i (*A. nigrofuscus*), maiko (*A. nigroris*), na'ena'e (*A. olivaceus*), manini (*A. triostegus*), pualu (*A. xanthopterus* and *A. blochii*), kole (*Ctenochaetus strigosus*), kala lolo (*Naso brevirostris*), kala holo (*N. hexacanthus*), umaumalei (*N. lituratus*), kala (*N. unicornis*), paki'i (*Bothus pantherinus*), humuhumu ele'ele (*Melichthys niger*), humuhumu hi'ukole (*M. vidua*) and the loulou (*Aluterus scriptus*). Besides these species as adults, juveniles of these and other species (e.g., the mu - *Monotaxis grandoculis*) were seen. Many of the adult individual fishes in the highly sought-after group were seen at varying distances away from the actual census areas, thus some species do not appear in the station counts (Appendix 1 of this and earlier reports).

Usually the most consistent location for finding many of the sought-after fish species both as adults and as juveniles is on the armor rock and steel protective cover for the Kahe plant warm-water discharge (Station 16). Because of the high degree of shelter provided by the armor rock as well as the well-developed coral community present on it and also due to the outfall (discharge), many species congregate there. Among these are many mamu (two species recognized, the Hawaiian mamu - *Abudefduf abdominalis* and the recently recognized species *Abudefduf vaigiensis*). Under the cover of the rocks are seen menpachi (*Myripristes amaneus*), aweoweo (*Priacanthus cruentatus*) and 'upapalu (*Apogon kallopterus*). In the December 2007 survey an estimated 200 grey mullet or ama'ama (*Mugil cephalus*) were encountered at Station 13. These fish had an average estimated length of 33 cm (~13 inches) contributing an estimated weight of 97.7 kilograms (215 lbs) to the standing crop at this station.

Many coral reef species other than fish are caught and consumed by people; among these are

specific algae and a number of invertebrates. Some individuals are interested in the collection of shells and when these usually cryptic species are seen at a station, they are so noted. Two species of molluscs have been seen on several occasions in the 2007-2013 surveys; these are the tiger cowry (*Cypraea tigris*) and the triton shell (*Charonia tritonis*). A species important in the making of fishing lures is the pearl oyster or pa (*Pinctado margaritifera*) which is protected by law and is commonly seen at many of the survey sites. The octopus or he'e (*Octopus cyanea*) was occasionally encountered at some of the stations. Individual he'e ranged from less than a pound in weight up to an estimated four pounds. The sought-after alga, limu kohu (*Asparagopsis taxiformis*) is seasonally common at many of the stations sampled in this study.

6. Standing Crops

Coral reefs function as relatively closed systems and thus in the pristine situation may represent the accumulation of carbon over a considerable period of time (Johannes *et al.* 1972). Some of this carbon is tied up in the living biomass of the reef of which fishes are only a part. Goldman and Talbot (1975) have suggested that a reasonable maximum biomass of coral reef fishes is approximately 200 g/m² or 2,000 kilograms per hectare (kg/ha). Space and cover are important agents governing the distribution of coral reef fishes (Sale 1977). Similarly the standing crop of fishes on a reef is correlated with the degree of vertical relief of the substratum (Risk 1972). Studies conducted on coral reefs in Hawai'i and elsewhere have estimated fish standing crops to range from 20 to 200 g/m² (Brock 1954, Goldman and Talbot 1975, Brock *et al.* 1979). Eliminating the direct impact of man due to fishing pressure and/or pollution, the variation in standing crop appears to be related to the variation in the local topographical complexity of the substratum which is governed, in part, by the degree of development in the coral community. Thus habitats with high structural complexity affording considerable shelter space usually harbor a greater estimated standing crop of coral reef fish; conversely, transects conducted in structurally simple habitats (e.g., sand flats) usually result in lower estimated standing crops (0.2 to 20 g/m²). Local studies (Brock and Norris 1989) suggest that with the manipulation (increasing) of habitat space or food resources (Brock 1987), fish standing crops may approach 2,000 g/m². Thus under certain circumstances, coral reefs may be able to support much larger standing crops of fishes than previously realized.

High standing crops (i.e., above 200 g/m²) were encountered during most surveys at several stations. In the 27 December 2007 survey at Station 9 where the estimated standing crop was 290 g/m², the opelu (*Decapterus macarellus*) made up 89% of this total at that location. Opelu are a coastal neritic species meaning that they school and move freely through the coastal waters which is very different than many coral reef fish species that have much smaller areas in which they forage. Similarly at Station 13 where the standing crop was estimated to be 594 g/m², the school of grey mullet or ama'ama (*Mugil cephalus*) described above comprised 82% of the total biomass. Again, ama'ama are usually seasonal in their appearance in coastal waters and travel over large areas of Hawai'i's waters. In the 4 April 2008 survey at Station 2, a school of 60 adult na'ena'e (*Acanthurus olivaceus*) swam through the census area bringing the total estimated biomass to 238 g/m² and these fish comprised 84% of the total weight present at this station. The

30 May 2008 survey noted a high standing crop at Station 16 (358 g/m²) where the mamo (*Abudefduf abdominalis* and *A. viagiensis*) made up 29% of the total and the kala lolo (*Naso brevirostris*) added 13% to the total estimated weight at this station. On 19 August 2008 at Station 16 the estimated biomass was 396 g/m² and again, the mamo comprised 51% of the total and a school of opelu passed through the census area and contributed 22% to the standing crop present at this station. In the 25 November 2008 survey at Station 16 where the estimated standing crop was 225 g/m², the two mamo species again comprised 38% of the biomass present at that time. In March 2009 survey Station 6 had an estimated standing crop of 259 g/m² and the palukaluka (*Scarus rubroviolaceus*) contributed 16% of the standing crop while the na'ena'e (*Acanthurus olivaceus*) added 40% to the biomass at this station. At Station 16 where the standing crop was estimated to be 577 g/m², two mamo species made up 15%, the kala lolo (*Naso brevirostris*) contributed 15% and the uhu (*Scarus sordidus*) added 22% to the standing crop present. In the May 2009 survey Station 5 had an estimated standing crop of 224 g/m² and the na'ena'e (*Acanthurus olivaceus*) made up 33% of it while at Station 16 where the standing crop was estimated to be 425 g/m², the two mamo species comprised 20% of the total weight present. The July 2009 survey noted that the estimated standing crop at Station 4 was 209 g/m² and the na'ena'e (*Acanthurus olivaceus*) made up 70% of it while at Station 5 the standing crop was 267 g/m² and again, na'ena'e made up 30% of the total biomass present. The standing crop of fishes at Station 16 was estimated to be 431 g/m² and the two mamo species made up 27% of it while the kala lolo (*Naso brevirostris*) added 8% to the biomass present.

The 29 March 2010 survey noted only one station with an estimated standing crop greater than 200 g/m²; this was Station 16 where the standing crop was 561 g/m² and the opelu (*Decapterus macarellus*) comprised 26% and the kala lolo (*Naso brevirostris*) made up 24% of the total estimated standing crop at this station. The 14 May survey encountered estimated standing crops in excess of 200 g/m² at two stations; Station 5 noted a biomass of 242 g/m² with the na'ena'e (*Acanthurus olivaceus*) contributing 62% of this estimated weight and the kole (*Ctenochaetus strigosus*) adding another 13% to the total weight at this station. The estimated standing crop at Station 16 was 390 g/m² and the two mamo species (*Abudefduf abdominalis* and *A. viagiensis*) contributed 35% of this estimated weight and the kala lolo (*Naso brevirostris*) added 25% to the total at this station. In the 12 August 2010 survey the standing crop at Station 15 was estimated to be 207 g/m² and the whitebar surgeonfish or maiko'iko (*Acanthurus leucoparicus*) provided 27% of this total and the na'ena'e (*Acanthurus olivaceus*) added 52% to the standing crop at this station. Again the estimated biomass at Station 16 in the August 2010 survey was elevated (603 g/m²) and five species were important contributors: the opelu (*Decapterus macarellus* - 25%), the two mamo species (*Abudefduf abdominalis* and *A. viagiensis* - 13%), the hinalea lauili (*Thalassoma duperrey* - 18%) and the kala lolo (*Naso brevirostris* - 11%). The 29 October 2010 survey found three stations with estimated standing crops in excess of 200 g/m². The standing crop at Station 7 was estimated to be 245 g/m² and the humuhumu ele'ele (*Melichthys niger*) added 49% to the total and the red weke or weke'ula (*Mulloidichthys vanicolensis*) contributed 26%. At Station 9 the standing crop was estimated to be 730 g/m² where a large school of white weke (*Mulloidichthys flavolineatus*) in the transect area made up 99% of the biomass at this station. Finally at Station 16 the biomass was estimated to be 554

g/m² and several schools of opelu (*Decapterus macarellus*) made up 41% of the total and a school of blue-lined snapper or ta'ape (*Lutjanus kasmira*) added 23% to the total at this station.

The 25 February 2011 survey noted only one station with a high estimated standing crop which was Station 16 on the Kahe discharge pipe where the estimated standing crop was 430 g/m² and the bluelined snapper or ta'ape (*Lutjanus kasmira*) comprised 52 percent of it and the kala lolo (*Naso brevirostris*) made up 21 percent of the estimated weight at this station. The 16 June survey found a high standing crop (273 g/m²) at Station 3 where the na'ena'e (*Acanthurus olivaceus*) made up 75 percent of the total estimated weight recorded at this station. At Station 6 the standing crop was estimated to be 206 g/m² and the palukaluka (*Scarus rubroviolaceus*) contributed 38 percent to this and the na'ena'e (*Acanthurus olivaceus*) added 27 percent to this estimated biomass while at Station 16 the standing crop was estimated to be 318 g/m² and the two mamo species (*Abudefduf abdominalis* and *A. viagiensis*) added 25 percent while the kala lolo (*Naso brevirostris*) contributed 33 percent to the total at this station. In the 29 July 2011 survey the standing crop at Station 3 was 435 g/m² and the na'ena'e (*Acanthurus olivaceus*) made up 89 percent of the total while at Station 6 where the standing crop was 234 g/m², the na'ena'e comprised 22 percent and the palukaluka (*Scarus rubroviolaceus*) added 24 percent to the biomass at this station. Again the standing crop at Station 16 was high (436 g/m²) and the ta'ape (*Lutjanus kasmira*) contributed 32 percent, the weke'ula (*Mulloidichthys vanicolensis*) added 23 percent while the kala lolo (*Naso brevirostris*) comprised 17 percent of the biomass at this station. Finally the 23 November 2011 survey found a high standing crop at Station 2 (263 g/m²) and Station 3 (379 g/m²) both due to the na'ena'e (*Acanthurus olivaceus*) making up 67 percent at Station 2 and 60 percent at Station 3. At Station 10 on this date, a school of weke'ula (*Mulloidichthys vanicolensis*) made up 71 percent of the estimated 318 g/m² and at Station 16 another school of weke'ula comprised 66 percent of the total 681 g/m² estimated biomass at this station.

In the four 2012 surveys, all standing crop estimates at stations in the 2 May and 23 July surveys were less than 200 g/m². However in the 23 May 2012 survey at Station 16 the standing crop of fishes was estimated to be 214 g/m² and the species contributing heavily to this biomass include the Indo-Pacific sergeant (*Abudefduf vaigiensis*) which added 19% to the station total. In the 2 November 2012 survey, the standing crop again at Station 16 was estimated to be 334 g/m² and a school of mackerel scad or opelu (*Decapterus macarellus*) contributed 50% to this biomass and the hinalea lauili (*Thalassoma duperrey*) added 14% and the Indo-Pacific sergeant (*Abudefduf vaigiensis*) accounted for 11% of the total at this station.

Four surveys were completed in 2013 with the first being carried out on 3 May. Standing crop estimates at three stations exceeded 200 g/m²; at Station 5 (Ko'Olina 1) the estimated biomass was 238 g/m² due largely to the manini (*Acanthurus triostegus*) making up 36% of the total and the na'ena'e (*Acanthurus olivaceus*) adding 46% to the total at this station. At Station 8 (Kahe 5-B) the standing crop was estimated to be 225 g/m² and the na'ena'e (*Acanthurus olivaceus*) made up 91% of the total at this station. Once again opelu (*Decapterus macarellus*) were present around the Kahe plant outfall (Station 16) during the 3 May survey comprising 50% of the total

standing crop that was estimated to be 348 g/m². The 14 June 2013 survey noted a high standing crop at Station 2 (238 g/m²) which was due to the na'ena'e (*Acanthurus olivaceus*) making up 85% of the estimated biomass at this station and at Station 16 (Kahe outfall) the standing crop was estimated to be 294 g/m² and the uhu (*Scarus sordidus*) comprised 27% of the total and the two mamo species (*Abudefduf abdominalis* and *A. vaigiensis*) contributed 21% to the total at this station. The 20 September 2013 survey noted an extremely high standing crop (1,619 g/m²) at Station 9 (Kahe 7B), 98% of which was due to a school of pelagic halfbeaks or iheihe (*Hyporhamphus pacificus*) passing through the transect area during the census. The standing crop at Station 12 (Kahe 7E) was estimated to be 372 g/m² and a passing school of the opelu (*Decapterus maracellus*) made up 75% of that total. Once again the standing crop at Station 16 (Kahe discharge) was estimated to be 413 g/m² and the opelu (*Decapterus macarellus*) contributed 11%, the uhu (*Scarus psittacus*) added 28% and the two mamo species (*Abudefduf abdominalis* and *A. vaigiensis*) provided 24% to the total for this station. The fourth quarter survey was carried out on 18 December 2013 where at Station 2 (CIP 2) the standing crop was estimated to be 281 g/m² and na'ena'e (*Acanthurus olivaceus*) added 67% and at Station 16 (Kahe outfall) the standing crop was estimated to be 296 g/m² and the weke'ula (*Mulloidichthys vanicolensis*) contributed 39%, the hinalea lauili (*Thalassoma duperrey*) added 15% and the two mamo species (*Abudefduf abdominalis* and *A. vaigiensis*) provided 13% to the total at this station.

A simple review of the above data finds that the same species often contribute substantially to the estimated standing crops at the same stations over time. A major reason for this is despite many stations not having a high degree of topographic complexity present in the 4 x 50 m transect areas, is the foraging behavior of the species involved. Since many of the fishes encountered on a transect such as the na'ena'e (*Acanthurus olivaceus*), uhus (*Scarus sordidus* and *S. psittacus*) and the palukaluka (*Scarus rubroviolaceus*) move and forage over areas considerably larger than the census area, they often will pass through the census area while it is being conducted and are thus counted. In other instances such as at the Kahe outfall station (Station 16), the presence of a unidirectional flow of warm discharge water containing particles that may serve as food as well as the high degree of topographical complexity all serve to draw both sedentary and more mobile fish species to the area including opelu (*Decapterus macarellus*) and weke'ula (*Mulloidichthys vanicolensis*).

7. Comparative Analysis of Early Hawaiian Electric Biological Data to the 2007-2013 Data

As noted above, Hawaiian Electric's environmental monitoring program for the Kahe Generating facility started in the 1970's, monitoring many of the same locations that are monitored today fronting the plant. These data are given in Coles *et al.* (1985b) and in a summary table (Table 33) in Coles *et al.* (1985a). Fish transect data from seven stations sampled in the 1976-1984 period fronting the Kahe Generating facility have been compared to the 2007-2013 data collected from those same sites. The sites include Station 7 (old #1-D started in 1979), Station 8 (old #5-B started in 1976), Station 10 (old #7-C started in 1976), Station 11 (old #7-D

started in 1976), Station 12 (old #7-E started in 1980), Station 13 (old #10-C started in 1979) and Station 14 (old Nanakuli control started in 1979). In this analysis, the annual means for the number of fish species and number of fish individuals encountered over those seven stations in common between the two groups of surveys are compared by addressing the question, “Are there any statistically significant differences among the annual mean number of fish species or annual mean number of individual fish censused per transect over the 1979-1984 and 2007-2013 periods?” Again, to address this question two non-parametric tests were used: the Kruskal-Wallis analysis of variance (ANOVA) and the Student-Newman-Kuels (SNK) Test where the Kruskal-Wallis ANOVA is used to demonstrate statistically significant differences among parameter means (by date) but cannot show where those differences are and the SNK Test is used to group related sample means and separate those means that are significantly different from one another.

The results of these analyses are given in Table 7 and referring to this table, it is shown that there are no significant differences among either of the annual means for the number of fish species seen per transect or the number of individual fish censused per transect despite the imposition of three major storm events. With respect to the annual mean number of species seen per transect, we find the greatest annual means occurring prior to the January 1980 storm event and the lowest mean (1983) occurring following Hurricane Iwa in 1982. With the annual mean number of individual fish seen per transect, the highest means occur with the recent (2007-2013) surveys and the lowest following Hurricane Iwa (1983) but the order among the dates does not parallel that for the mean number of fish species (Table 7). Thus not all species of fish were impacted to the same degree with the occurrence of these two early high wave events. Fish standing crop information was not readily available for Stations 7, 8, 10, 11, 12, 13 or 14 in the early (1976-1984) Hawaiian Electric dataset except for 1984 thus was not included in the above (Table 7) analysis. However, the non-parametric Wilcoxon Two-Sample Test was used to examine the mean estimated standing crop of fishes in 1984 at the above seven stations comparing this mean to the mean estimated biomass at these stations in the 2007-2013 dataset. Despite the mean estimated standing crop (here 53 g/m²) being greater in 2007-2013 than in 1984 (26 g/m²), the Wilcoxon Two-Sample Test failed to find any statistically significant differences ($p > 0.26$, n.s., where a $p < 0.05$ signifies significance) in the estimated standing crop at these seven stations sampled minimally 22 years apart. Again the standing crop statistical results support those found with the mean number of fish species or the mean number of individual fish censused per transect (Table 7).

In summary, there are no statistically significant differences among the annual mean number of fish species or individuals censused utilizing data that span a 37-year period (1976-2013) at seven monitoring stations fronting the Kahe Generating Station despite the imposition of three major storm events in 1980, 1982 and 1992 (see Section 2 of this report). These data suggest that the fish communities have to some extent recovered from these disturbances.

8. Federally Protected Species

When encountered during field work, federally protected species are noted. Five species that are encountered (or heard underwater) around the high Hawaiian Islands are the green turtle (*Chelonia mydas*), the hawksbill turtle (*Eretmochelys imbricata*), the spinner porpoise (*Stenella longirostris*), the Hawaiian monk seal (*Monachus schauinslandi*) and present seasonally, the humpback whale (*Megaptera novaeangliae*).

Because of low population numbers, the Hawaiian green sea turtle was given protection under the federal Endangered Species Act in the mid-1970's. Green turtles as adults are known to forage and rest in the shallow waters around the main Hawaiian Islands. Reproduction in the Hawaiian population occurs primarily during the summer months in the Northwest Hawaiian Islands with adults migrating during the early summer to these isolated atolls and returning in the late summer or early fall. In the main Hawaiian Islands, green turtles rest along ledges, caves or around large coral mounds in coastal waters usually from 15 to 20 m in depth during the day. Under the cover of darkness, turtles will travel inshore to shallow subtidal and intertidal habitats for foraging on algae or limu. (Balazs *et al.* 1987). The normal range of these daily movements between resting and foraging areas is about one kilometer (Balazs 1980, Balazs *et al.* 1987). In general, appropriate algal forage for these turtles is found in shallow waters inshore of the resting areas. Selectivity of algal species consumed by Hawaiian green turtles appears to vary with the locality of sampling, but stomach content data show *Acanthophora spicifera* (an introduced species) and *Amansia glomerata* to quantitatively be the most important (Balazs *et al.* 1987); the preferences may be due to the ubiquitous distribution of these algal species.

The Hawaiian green turtle population has rebounded under the more than 30 years of federal protection afforded to it such that today, green turtles are commonly seen in the waters fronting most beaches around the islands. In contrast, the hawksbill turtle is much less common and much less is known about its biology in Hawaiian waters. Hawksbill turtles do not attain the size of green turtles in Hawaiian waters, nest on very small and isolated beaches around the main islands and are omnivorous in their feeding habitats. In the waters surveyed under the present study, no hawksbill turtles have been seen by us.

Green turtles were observed on fourteen of the twenty-four surveys completed to date. All turtles seen were juveniles (i.e., having a carapace length estimated to be less than 75 cm) except for a pair of adults (estimated straight-line carapace lengths = 90 cm) in the 10 August 2010 survey at Station 9 and a single individual (~85 cm) at Station 5 in the 20 September 2013 survey. Some turtles were sleeping while others observed were actively swimming. There is a depression in the limestone at Station 8 where green turtles often rest; in 2009 a small (~45 cm straight-line carapace length) green turtle was observed in this depression in the 19 March and 21 July 2009 surveys. In the August 2010 and 16 June 2011 surveys this same depression was occupied by a ~65 cm sleeping juvenile turtle. On the 23 November 2011 survey two turtles (~60 cm and ~70 cm) were found in this same depression sleeping and on the 23 May 2012 survey this depression was occupied by an ~75 cm resting turtle. In the 14 June 2013 survey an

approximate 45 cm turtle was observed resting in the same depression at station 8 and in the 20 September 2013 survey a 60 cm turtle was resting at this same location. This same depression was once again occupied by a ~40 cm turtle in the 18 December 2013 survey. In the 25 November 2008 survey six green turtles were found resting on the bottom in a depression just seaward of Station 5 and again on the 23 May 2012 survey a single juvenile turtle (~70 cm) was seen at Station 16 (the Kahe Generating Station warm-water discharge) swimming in a northwest direction and in the 18 December 2013 survey a 40 cm individual green turtle was seen swimming towards shore along the discharge pipe. In the 14 June 2013 survey, green turtles were also seen at Station 6 (~70 cm) and a second individual at 9 (~75 cm). In no cases were any tags or tumors seen on any of the turtles sighted to date.

For many years, Hawaiian monk seals were not seen very often around the main Hawaiian Islands but the population was located in the Northwest Hawaiian Islands and over time the population numbers have declined. Despite this, in recent years an increasing number of Hawaiian monk seals are now seen on the beaches around the main islands with the occasional female giving birth on island beaches. The reason(s) for these changes in the population are unknown but the result is monk seals are now occasionally seen while carrying out environmental surveys around the main islands. On the 30 May 2008 survey an adult male monk seal approached the vessel while at anchor at Station 14. This seal carried a tag (not readable at distance) and it swam around the vessel and subsequently left heading towards the shoreline. This seal has not been seen since. In the 14 June 2013 survey while conducting census work on the Kahe facility discharge pipe, a spear fisherman swam up to the survey vessel and reported that two monk seals were presently on the beach resting near the discharge pipe. He said that on the day prior, this pair of seals took all of his speared fish and one of the seals was “aggressive”. These seals were not observed by us.

It should be noted that the endangered humpback whale is known to frequent island waters in their annual migrations to Hawaiian wintering grounds. They normally arrive in island waters about December and depart by April. In general their distribution in Hawaii appears to be limited to the 180 m (100 fathom) isobath and in shallower waters (Nitta and Naughton 1989). Whales were observed further seaward of the Barbers Point - Kahe Point study area and their songs could be heard underwater during the 27 December 2007, the 19 March 2009, the 24 March 2010, the 25 February 2011 and 18 December 2013 surveys.

Spinner porpoises are occasionally seen in the Kahe Point area and were first encountered there during this study on the 30 May 2008 survey where three pods were seen each having about 35 individuals present. In the 14 May 2010 survey a pod of about 30 individuals passed by during a census at station 10 and in the 23 July 2012 survey a small pod of approximately 20 porpoises were seen traveling northwest seaward of Stations 5 and 6. On the 18 December 2013 survey a small pod of spinner porpoises (estimated 15 individuals), were seen moving west just seaward of Station 7. Hawaiian spinner porpoises are known to rest in shallow bays during the day and at night move offshore to feed on midwater fishes and squids that rise to the surface to forage.

9. Long-Term Perspective on the Barbers Point-Kahe Point Fish Communities

As noted and documented above, the three early storm events (1980, 1982 and 1992) all impacted marine communities offshore of the Barbers Point - Kahe Point areas. These impacts were probably greatest on the coral communities which due to their sessile nature, must withstand the wave forces impinging on them or perish. Corals are relatively slow-growing and depending on the species, individual colonies may live for a considerable time and in doing so create habitat for fishes and other reef species. If disturbance to the coral community is relatively frequent, surviving corals probably do not contribute much to the three-dimensional structure of the habitat, thus keeping the fish community development in an earlier successional stage than it might otherwise be. Storms not only directly impact the living resources but also the geological condition of reef areas. As noted by the early Hawaiian Electric studies, considerable sand movement occurred with the first two major storms that occurred in 1980 and 1982 and today much of the area west of the Kahe facility's ocean outfall is now nearly devoid of sand leaving a near-featureless hard bottom that is scoured with passing small wave events that retard benthic and fish community development. A similar situation exists east of the Barbers Point Harbor entrance channel where considerable hard (limestone) substratum is present with much of it having poor benthic community development. This again results in a poorly developed resident fish community, which is what we see in much of the area today and did so forty years ago (Brock, personal observations). Thus the measures of fish community development used here (the diversity of species and numbers of individuals present as well as the standing crop) do not suggest well-developed resident fish communities at many of the sample sites. However where topographical complexity is greater and benthic communities are better developed, the resulting fish communities are well-developed. This is best illustrated at Station 16 located at the terminus of the Kahe facility discharge, where, despite high use which includes snorkel/dive tours as well as spear fishermen, the fish community remains relatively well-developed. The high degree of development in the resident fish communities on the Kahe discharge structure lend further support to the lack of negative impact due to the operation of the discharge.

Given the long-term extant data set spanning 37 years and the apparent lack of strong significant changes occurring with the three early (1980, 1982 and 1992) storm events (which is probably due to some level of recovery in the intervening period), suggests that the variation seen in the measures of the fish community used here will continue to fluctuate at a similar magnitude in future monitoring events as this program moves forward. The 37 years of well-documented environmental history for the Barbers Point - Kahe Point area (completed largely by the Hawaiian Electric environmental program), provides much of the explanation to the degree of development of resident fish communities we encounter in the area today.

LITERATURE CITED

Bak, R. and B. Luckhurst. 1980. Constancy and change in coral reef habitats along depth gradients at Curacao. *Oecologia* 47:145-155.

- Balazs, G.H. 1980. Synopsis of biological data on the green turtle in the Hawaiian Islands. NOAA Tech. Memorandum, NMFS. NOAA-TM-NMFS-SWFC-7. 141p.
- Balazs, G.H., R.G. Forsyth and A.K.H. Kam. 1987. Preliminary assessment of habitat utilization by Hawaiian green turtles in their resident foraging pastures. NOAA Tech. Memorandum, NMFS. NOAA-TM-NMFS-SWFC-71. 107p.
- Ball, M., E. Shinn and K. Stockman. 1967. The geological effects of Hurricane Donna in south Florida. J. Geol. 75:583-597.
- Blumenstock, D., F. Fosberg and C. Johnson. 1961. The re-survey of typhoon effects on Jaluit Atoll in the Marshall Islands. Nature 189:618-620.
- Brock, R.E. 1982. A critique on the visual census method for assessing coral reef fish populations. Bull. Mar. Sci. 32:269-276.
- Brock, R.E. 1987. An assessment of the fishes and macrobenthos off the Kahuku coast, Oahu in relation to an aquaculture effluent. Prepared for AECOS, Inc., Kailua, Hawaii. 16p.
- Brock, R.E. 1992a. Status of coral communities fronting the Kahe Generation Facility, Kahe Point, Oahu following Hurricane Iniki. Prepared for the Environmental Department, Hawaiian Electric Co., Inc., Honolulu. Environmental Assessment Co. Report No. 92-17. 12p.
- Brock, R.E. 1992b. Status of the green turtle (*Chelonia mydas*) population fronting the proposed Campbell drainage channel to serve the Kapolei Business Park. Prepared for Engineering Concepts, Inc., 250 Ward Avenue, Suite 206, Honolulu. Environmental Assessment Co. Report No. 92-15. 14p.
- Brock, R.E. 1996. A study of the impact of Hurricane Iniki on coral communities at selected sites in Mamala Bay, O'ahu, Hawai'i. Project Report PR-96-09. Water Resources Research Center, University of Hawaii, Honolulu. vii+23p.
- Brock, R.E. 2009. CIP generation project 2008 community benefits program: reef fish monitoring project year 1 results. Prepared for the Environmental Department, Hawaiian Electric Co., Inc., P.O. Box 2750, Honolulu, Hawaii 96840-0001. Environmental Assessment, LLC, Report No. 2009-07. 38p+appendices.
- Brock, R.E. 2010. CIP generation project 2009 community benefits program: reef fish monitoring project year 2 results. Prepared for the Environmental Department, Hawaiian Electric Co., Inc., P.O. Box 2750, Honolulu, Hawaii 96840-0001. Environmental Assessment, LLC, Report No. 2010-009. 38p+appendices.
- Brock, R.E. 2011. CIP generation project 2010 community benefits program: reef fish

monitoring project year 3 results. Prepared for the Environmental Department, Hawaiian Electric Co., Inc., P.O. Box 2750, Honolulu, Hawaii 96840-0001. Environmental Assessment, LLC, Report No. 2011-002. 41p+appendices.

Brock, R.E. 2012. CIP generation project 2011 community benefits program: reef fish monitoring project year 4 results. Prepared for the Environmental Department, Hawaiian Electric Co., Inc., P.O. Box 2750, Honolulu, Hawaii 96840-0001. Environmental Assessment, LLC, Report No. 2012-010. 42p+appendices.

Brock, R.E. 2013. CIP generation project 2012 community benefits program: reef fish monitoring project year 5 results. Prepared for the Environmental Department, Hawaiian Electric, P.O. Box 2750, Honolulu, Hawaii 96840-0001. Environmental Assessment, LLC, Report No. 2013-008, 45p+appendices.

Brock, R.E., C. Lewis and R.C. Wass. 1979. Stability and structure of a fish community on a coral patch reef in Hawaii. *Mar. Biol.* 54:281-292.

Brock, R.E. and J.E. Norris. 1989. An analysis of the efficacy of artificial reef designs in tropical waters. *Bull. Mar. Sci.* 44:934-941.

Brock, V.E. 1954. A preliminary report on a method of estimating reef fish populations. *J. Wildlife Mgmt.* 18:297-308.

Coles, S.L. 1979. Annual report. Kahe Generating Station. NPDES monitoring program, vol. 1 text. Honolulu. 263pp

Coles, S.L. and D.T. Fukuda. 1984. Annual report: Kahe generating station NPDES monitoring program. Vol. I, Text. Environmental Department, Hawaiian Electric Co., Inc., P.O. Box 2750, Honolulu, Hawaii 96840.

Coles, S.L., D.T. Fukuda and C.R. Lewis. 1981. Annual report. Kahe Generating Station. NPDES monitoring program, vol.1 text. Honolulu, 237 pp.

Coles, S.L., D.T. Fukuda and C.R. Lewis. 1982 Annual report. Kahe Generating Station. NPDES monitoring program, vol.1 text. Honolulu, 221p.

Coles, S.L., D.T. Fukuda and K.S. Oda. 1985a. Annual report Kahe Generating Station NPDES monitoring program. Vol. I. Text. Environmental Department, Hawaiian Electric Company, Inc., P.O. Box 2750, Honolulu, Hawaii 96840. v+172p.

Coles, S.L., D.T. Fukuda and K.S. Oda. 1985b. Annual report. Kahe generating station NPDES monitoring program. Vol. II, Appendices. Environmental Department, Hawaiian Electric Company, Inc., P.O. Box 2750, Honolulu, Hawaii 96840, 168p.

- Connell, J. 1978. Diversity in tropical rain forests and coral reefs. *Science* 199:1302-1310.
- Cooper, M. 1966. Destruction of marine fauna and flora in Fiji caused by the hurricane of February 1965. *Pacif. Sci.* 20:137-141.
- Dollar, S. 1982. Wave stress and coral community structure in Hawaii. *Coral Reefs* 1:71-81.
- Dollar, S. and G. Tribble. 1993. Recurrent storm disturbance and recovery: a long-term study of coral communities in Hawaii. *Coral Reefs* 12:223-233.
- Done, T., P. Dayton, A. Dayton and R. Steger. 1991. Regional and local variability in recovery of shallow coral communities: Moorea, French Polynesia and central Great Barrier Reef. *Coral Reefs* 9:183-192.
- Goldman, B. and F.H. Talbot. 1975. Aspects of the ecology of coral reef fishes. In: *Biology and Geology of Coral Reefs*, ed. O.A. Jones and R. Endean, pp. 125-154. Vol. III, Biology 2. Academic Press, New York.
- Grassle, J. 1973. Variety in coral reef communities. *Biology and geology of coral reefs*, Biology 1, vol. 2, ed. O.A. Jones and E. Endean, pp. 247-270. New York: Academic Press.
- Grigg, R.W. 1983. Community structure, succession and development of coral reefs in Hawaii. *Mar. Ecol. Prog. Ser.* 11:1-14.
- Grigg, R.W. and J.E. Maragos. 1974. Recolonization of hermatypic corals on submerged lava flows in Hawaii. *Ecology* 55:387-395.
- Harmelin-Vivien, M. and P. Laboute. 1986. Catastrophic impact of hurricanes on atoll outer reef slopes in the Tuamotu (French Polynesia). *Coral Reefs* 5:55-62.
- Hiatt, R.W. and D.W. Strasburg. 1960. Ecological relationships of the fish fauna on coral reefs of the Marshall Islands. *Ecol. Monogr.* 30:65-127.
- Hobson, E.S. 1974. Feeding relationships of teleostean fishes on coral reefs in Kona, Hawaii. *Fish. Bull.* 72:915-1031.
- Johannes, R.E., *et al.* 1972. The metabolism of some coral reef communities: a team study of nutrient and energy flux at Eniwetok. *Bioscience* 22:541-543.
- Knutson, D.W., R.W. Buddemeier and S.V. Smith. 1972. Coral chronometers: seasonal growth bands in reef corals. *Science* 177:270-272.
- Maragos, J., G. Baines and P. Beveridge. 1973. Tropical cyclone Bebe creates a new land

formation on Funafuti Atoll. *Science* 181:1161-1164.

Nitta, E.T. and J.J. Naughton. 1989. Species profiles: life histories and environmental requirements of coastal vertebrates and invertebrates, Pacific Ocean region; Report No. 2, humpback whale, *Megaptera novaeangliae*. Tech. Rept. EL-89-10. Prepared by NMFS, NOAA, Honolulu for the U.S. Army Engineer Waterways Experiment Station, Ms.

Noda, E.K. 1983. Affects of Hurricane Iwa, November 23, 1982 offshore of Kahe Point, Oahu. Unpublished report prepared for Research Corp. of the University of Hawaii.

Ogg, J. and J. Koslow. 1978. The impact of typhoon Pamela (1976) on Guam's coral reefs and beaches. *Pacif. Sci.* 32:1056-118.

Perkins, R. and P. Enos. 1968. Hurricane Betsy in the Florida-Bahamas area - Geological effects and comparison with Hurricane Donna. *J. Geol.* 76:710-717.

Randall, J.E. 2007. Reef and shore fishes of the Hawaiian Islands. Sea Grant College Program, University of Hawai'i, Honolulu. xiv+546p.

Ricker, W.E. 1975. Handbook of computations for biological statistics of fish populations. Fish. Res. Bd. Canada, Bull. 119. 300p.

Risk, M.J. 1972. Fish diversity on a coral reef in the Virgin Islands. *Atoll. Res. Bull.* 153:1-6.

Sale, P.F. Maintenance of high diversity in coral reef fish communities. *Amer. Nat.* 111:337-359.

Stoddart, D. 1969. Post-hurricane changes on the British Honduras reefs and cays: re-survey of 1965. *Atoll Res. Bull.* 13:1-25.

Stoddart, D. 1974. Post-hurricane changes on the British Honduras reefs: re-survey of 1972. *Proc. 2nd Int. Coral Reef Symp.* 2:473-483.

Veron, J.E.N. 1995. Corals in space and time. UNSW Press, Sydney, NSW, Australia 2052.

Walsh, W.J. 1983. Stability of a coral reef fish community following a catastrophic storm. *Coral Reefs* 2:49-63.

Woodley, J. *et al.* 1981. Hurricane Allen's impact on Jamaican coral reefs. *Science* 214:749-755.

TABLE 1. Latitude and Longitude waypoints (in decimal minutes) for each of the sixteen permanently marked fish monitoring stations utilized in this study (GPS waypoints courtesy of the Environmental Department, Hawaiian Electric). Note that the first survey carried out on 27 December 2007 did not sample station numbers 5, 6, 7 and 16. The second survey on 4 April 2008 missed station 16 while surveys carried out subsequently on 22 occasions have sampled all sixteen sites.

Station No.	Station Area Name	Latitude	Longitude	Remarks
1	East 1	21°18.237' N	158°07.024'W	New- offshore CIP
2	East 2	21°18.452'N	158°07.152'W	New - offshore CIP
3	East 3	21°18.558'N	158°07.239'W	New - offshore CIP
4	East 4	21°18.406'N	158°07.285'W	New - offshore CIP
5	Ko'Olina 1	21°19.724'N	158°07.581'W	New - offshore Ko'Olina
6	Ko'Olina 2	21°19.904'N	158°07.693'W	New - offshore Ko'Olina
7	HECO station 1D	21°20.763'N	158°07.773'W	Old Hawaiian Electricstation
8	HECO station 5B	21°21.145'N	158°07.819'W	Old Hawaiian Electric station
9	HECO station 7B	21°21.239'N	158°07.855'W	Old Hawaiian Electric station
10	HECO station 7C	21°21.255'N	158°07.881'W	Old Hawaiian Electric station
11	HECO station 7D	21°21.268'N	158°07.893'W	Old Hawaiian Electric station
12	HECO station 7E	21°21.272'N	158°07.977'W	Old Hawaiian Electric station
13	HECO station 10C	21°21.522'N	158°07.925'W	Old Hawaiian Electric station
14	Nanakuli Control 1	21°22.329'N	158°08.440'W	Old Hawaiian Electric station
15	Nanakuli Control 2	21°22.353'N	158°08.462'W	New control station
16	On Outfall	21°21.193'N	158°07.869'W	New north side of outfall

TABLE 2. Summary of the fish censuses carried out at sixteen locations on twenty-four surveys over the December 2007 - December 2013 period. The percent of the total biomass is that assigned to each of five trophic categories: herbivores, planktivores, omnivores, carnivores and coral feeders is also given. Note that these percentages are rounded to the nearest whole number.

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
27-Dec-07	1	12	69	15	18		1	51	30
	2	19	155	143	87			9	4
	3	30	189	41	28		6	51	15
	4	Not sampled							
	5	Not sampled							
	6	Not sampled							
	7	28	306	92	40		40	19	1
	8	25	241	43	51	7	3	39	
	9	23	259	290	6	1	1	92	
	10	17	261	154		9	3	88	
	11	13	23	104	6		5	82	7
	12	34	581	63	21	1	24	51	3
	13	31	580	594	85	3	1	11	
	14	18	124	7	23	2	3	72	
	15	23	164	94	51		8	40	1
	16	Not sampled							
04-Apr-08	1	10	129	8		1	1	59	39
	2	25	333	238	89		1	9	1
	3	18	146	21	38		7	54	1
	4	25	270	116	57		3	37	3
	5	34	307	146	81	2	2	13	2
	6	31	292	164	67	1	2	29	1
	7	21	365	158	14		75	11	
	8	27	499	29	26	5	4	64	1
	9	17	75	74	25	1	1	73	
	10	11	117	8	42	1	5	52	
	11	6	21	4		1	2	97	
	12	25	390	31		1	15	79	5
	13	16	401	62	3	15	7	70	5
	14	12	260	14	1	1		98	
	15	17	214	129	83		1	15	1
	16	Not sampled							
30-May-08	1	12	77	9		1	17	82	
	2	21	220	64	65			34	1
	3	22	136	37	24		9	62	4
	4	30	293	49	28	1	23	45	3
	5	30	250	84	73		8	20	2
	6	32	265	132	77	1	7	14	1
	7	24	292	94	21		53	25	1
	8	26	412	75	70	9	1	20	
	9	21	152	95	21	67	1	11	
	10	21	167	55	60	14	3	23	
	11	12	81	21	35	2	37	26	
	12	25	453	14		4	28	60	8
	13	24	263	24	5	11	18	66	
	14	26	188	20	9		1	67	23
	15	13	80	34	69		3	26	2
	16	42	1205	358	8	43	2	47	

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
19-Aug-08	1	19	155	13	1		9	90	
	2	20	280	120	85		2	13	
	3	23	231	40	27		5	66	2
	4	26	415	108	43	8	6	43	
	5	24	227	69	67		9	22	2
	6	35	302	165	79	1	6	14	
	7	24	213	65	9		56	35	
	8	27	463	39	49	1	2	47	
	9	23	235	34	56	4	6	34	
	10	39	201	33	9	1	5	85	
	11	32	126	41	1	2	23	57	17
	12	23	514	33	19	2	13	56	10
	13	21	385	63	45	16	4	35	
	14	19	192	8	4	1		95	
	15	15	104	16	44	1	2	47	6
	16	37	1023	396	3	55	1	41	
25-Nov-08	1	6	20	2			6	53	40
	2	10	41	4	21		6	73	
	3	21	100	12	47	3	3	46	1
	4	20	165	79	54		1	45	
	5	31	289	91	81		1	17	1
	6	36	263	189	82	2	4	10	2
	7	31	394	60	37		36	27	
	8	33	147	29	49	6	1	43	1
	9	25	374	171	14	1		85	
	10	31	364	62	45	4	2	49	
	11	9	52	18	44	1	2	53	
	12	31	426	19	17	6	30	38	9
	13	32	931	155	20	57	4	18	1
	14	19	170	15	38		1	61	
	15	24	234	171	91		2	7	
	16	40	1017	225	10	49	1	39	1
19-Mar-09	1	14	93	13	11		1	83	5
	2	14	102	15	16		2	79	3
	3	22	126	21	18		23	50	8
	4	18	125	25	21		18	61	
	5	27	302	113	82		2	14	2
	6	33	370	259	91	2	1	5	1
	7	32	349	91	41	1	44	13	1
	8	21	353	31	32	1	3	63	1
	9	17	111	74	6		2	92	
	10	13	52	14	35			64	1
	11	5	7	4		1		99	
	12	28	251	15	34	2	2	57	5
	13	30	458	84	17	5	6	72	
	14	17	84	7	35		2	63	
	15	23	148	115	92		1	6	1
	16	48	1438	577	31	34	2	32	1

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
11-May-09	1	11	108	12	22		1	77	
	2	18	231	41	27		1	68	4
	3	26	224	65	64		7	27	2
	4	25	328	61	58		3	36	3
	5	31	383	224	87		3	9	1
	6	30	240	153	86	2	4	6	2
	7	26	263	51	31	1	45	22	1
	8	27	363	35	56	4	9	30	1
	9	15	88	20	51		1	48	
	10	20	159	22	32	1	14	52	1
	11	4	9	12			7	93	
	12	24	267	20	13	1	11	74	1
	13	28	459	147	20	8	1	71	
	14	11	43	6	25		8	67	
	15	17	194	174	87		1	12	
	16	39	1333	425	35	22	6	37	
21-Jul-09	1	17	141	18	2		9	81	8
	2	25	389	73	52			43	4
	3	31	301	80	26	5	31	34	4
	4	27	506	209	80		4	15	1
	5	39	582	267	65	5	6	23	1
	6	37	354	188	74	2	7	16	1
	7	33	589	155	28	2	49	21	
	8	26	800	47	47	2	7	44	
	9	27	204	70	6	4	3	87	
	10	24	212	30	15	42	2	41	
	11	10	40	12		1	2	97	
	12	26	432	20	18	6	18	46	12
	13	24	405	145	7	11	1	81	
	14	15	111	9	1	1	2	96	
	15	21	258	140	77	6	7	8	2
	16	40	1605	431	5	36	3	56	
29-Mar-10	1	17	162	30	56	0	9	25	10
	2	22	315	33	34	0	4	57	5
	3	27	197	45	70	1	10	17	2
	4	24	324	105	65	0	2	32	1
	5	31	312	129	76	4	8	10	2
	6	29	313	176	85	1	5	9	1
	7	26	336	67	26	0	46	28	0
	8	29	265	56	51	2	5	42	
	9	19	83	18	23	2	0	74	1
	10	13	53	10	40	0	21	38	
	11	10	28	14	1	0	4	95	
	12	24	245	54	7	54	15	23	0
	13	34	312	69	18	14	6	62	0
	14	11	101	7	31	0	2	65	2
	15	24	149	77	75		4	20	1
	16	29	1192	561	24	27	2	47	

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
14-May-10	1	18	94	15	55	0	8	37	
	2	17	91	14	33		7	48	13
	3	23	160	63	70	0	3	24	2
	4	16	326	85	71	0	6	19	4
	5	35	511	242	87	3	4	6	1
	6	37	241	164	82	2	3	14	1
	7	23	395	113	11	1	39	49	1
	8	26	361	80	78	2	6	13	0
	9	28	179	159	24	3	1	72	0
	10	21	119	55	53	24	2	20	0
	11	9	43	21		1	36	63	
	12	25	299	51	31	18	26	23	1
	13	31	369	57	9	35	5	50	0
	14	10	19	2	22		8	70	
	15	26	201	139	91	0	1	8	0
	16	33	1767	390	13	63	8	16	
12-Aug-10	1	22	198	157	69	0	1	31	0
	2	25	313	69	34	0	6	59	1
	3	25	225	28	42	0	8	49	1
	4	22	358	151	67	0	12	21	
	5	36	426	163	73	1	7	19	1
	6	30	233	118	63	2	11	23	1
	7	26	271	100	29	0	40	31	1
	8	24	425	62	73	1	5	21	0
	9	28	104	40	47	7	0	46	0
	10	20	106	31	24	49	4	23	0
	11	13	58	19	9	2	36	53	
	12	31	317	24	29	31	15	25	0
	13	32	359	60	11	12	10	68	
	14	13	51	23	85	0	0	14	2
	15	26	248	207	89	0	1	10	0
	16	33	1584	603	14	27	1	58	0
29-Oct-10	1	14	104	96	79		0	21	
	2	13	208	56	73		0	26	0
	3	27	183	49	61	0	6	32	2
	4	22	195	66	61	0	8	31	
	5	38	315	98	69	0	5	24	2
	6	36	294	123	79	2	3	15	2
	7	31	743	245	7	0	50	42	0
	8	28	262	24	33	1	6	60	0
	9	22	467	730	0	0	0	99	0
	10	17	57	21	31	0	10	59	1
	11	13	38	15	1	1	34	64	
	12	36	334	23	34	2	13	50	1
	13	35	478	192	23	5	1	69	1
	14	9	57	7		0	1	99	
	15	28	169	31	24	0	11	46	19
	16	35	1039	554	7	16	1	76	

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
25-Feb-11	1	9	42	5	13		1	86	
	2	16	183	66	83		1	17	
	3	17	119	18	17		16	66	1
	4	20	266	25	47	1	18	34	
	5	31	307	99	54	0	4	40	1
	6	27	328	196	49	6	4	40	1
	7	18	235	93	8		67	25	
	8	25	307	33	13	7	2	77	0
	9	13	61	10	11	19	21	48	0
	10	7	26	4		0	19	80	
	11	8	15	12		0	41	59	
	12	24	243	14	29	7	21	42	1
	13	27	427	119	22	19	5	54	
	14	9	32	2	13	1	3	83	
	15	14	69	23	28	0	6	66	0
	16	24	910	430	8	32	2	59	0
16-Jun-11	1	18	162	124	91	0	1	8	0
	2	17	123	66	78	0	2	19	0
	3	27	275	273	88		1	10	1
	4	25	340	80	66	0	6	28	0
	5	24	270	74	63	5	9	22	1
	6	33	281	207	82	1	8	8	0
	7	27	434	131	35	0	32	32	
	8	27	464	37	60	6	6	28	
	9	15	54	14	25	0	18	56	0
	10	16	103	13	6	1	0	93	
	11	11	42	6	1	1	0	98	
	12	28	769	50	2	54	8	36	0
	13	29	383	75	3	8	8	79	3
	14	12	88	5	0	1		99	
	15	21	340	108	94	0	3	3	1
	16	40	1315	318	17	59	6	17	0
29-Jul-11	1	16	137	14	4		8	89	0
	2	21	183	52	59	0	2	39	0
	3	23	277	435	96	1	2	1	0
	4	26	299	52	42	0	13	44	
	5	34	333	138	88	0	1	10	1
	6	36	375	234	86	1	5	6	1
	7	23	309	100	8	2	55	35	
	8	33	802	38	42	12	3	39	4
	9	22	477	285	13	0	0	86	0
	10	11	58	5	0	1		98	
	11	9	53	2	1	9	1	90	
	12	32	297	22	3	2	42	53	1
	13	33	327	36	13	23	10	53	0
	14	12	67	5	5	0	1	93	
	15	22	113	82	84		5	11	
	16	38	864	436	4	25	1	70	0

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
23-Nov-11	1	15	179	161	92		0	8	0
	2	22	348	263	88		0	11	1
	3	38	320	379	92		2	5	1
	4	26	360	166	81	0	4	14	1
	5	29	320	122	83	0	3	13	0
	6	30	291	188	85	0	3	11	1
	7	26	244	68	16	0	64	19	1
	8	27	343	32	45	7	5	41	1
	9	23	102	29	61	0	4	34	0
	10	20	85	19	40	0	5	53	2
	11	13	26	50	5		10	86	
	12	34	691	24	30	4	21	44	1
	13	35	1253	318	1	6	2	91	
	14	12	44	7	56	0	1	43	
	15	17	85	16	56	0	17	21	6
	16	28	1318	681	10	19	0	70	0
02-May-12	1	9	74	16	6		7	87	
	2	13	130	27	59	0	0	40	1
	3	23	137	65	66	7	3	23	2
	4	26	251	128	52	5	3	41	
	5	29	227	93	73	0	3	22	1
	6	35	276	147	75	2	4	18	1
	7	25	315	82	17	0	43	40	0
	8	31	371	130	56	0	4	39	0
	9	21	116	20	32	1	12	54	1
	10	15	78	16	20	0	7	71	1
	11	11	31	67	79		0	20	0
	12	28	262	50	31	1	16	52	1
	13	35	339	173	7	4	3	85	0
	14	14	89	9	20	0	1	79	
	15	20	150	54	84	0	4	12	
	16	26	568	143	20	40	6	33	0
23-May-12	1	15	105	52	84	0	0	16	
	2	15	194	53	60	0	3	36	1
	3	23	176	75	69	0	4	26	2
	4	18	357	49	36	1	11	53	
	5	28	211	57	73	0	4	22	1
	6	32	259	163	85	1	6	8	0
	7	19	247	48	39	1	21	40	
	8	22	270	42	36	0	5	58	0
	9	17	59	20	25	0	5	70	0
	10	13	36	10	65		10	23	1
	11	9	23	30	44		0	56	
	12	18	211	27	9	1	32	57	1
	13	28	211	71	5	6	5	84	
	14	11	89	4	52	2	1	45	
	15	17	118	19	23	0	23	54	
	16	23	846	214	14	27	2	58	0

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
23-Jul-12	1	23	274	189	67	0	0	32	0
	2	18	187	55	63	0	2	35	0
	3	21	114	39	65		13	21	2
	4	19	344	36	16	1	15	68	
	5	30	185	46	60	1	10	26	3
	6	30	184	134	79	1	3	16	0
	7	24	249	50	26	0	54	20	0
	8	29	212	41	57	0	3	39	1
	9	25	81	26	43	0	6	51	1
	10	13	64	9	16	1	11	71	1
	11	9	20	82	1	2		97	
	12	24	274	32	5	1	35	55	4
	13	34	439	92	2	14	5	78	1
	14	15	54	6	1	0	1	97	
	15	19	102	88	80	11	3	5	1
	16	30	685	153	28	44	4	25	0
02-Nov-12	1	18	201	86	76	0	2	22	
	2	17	224	52	57	0	2	40	0
	3	24	147	109	86		2	11	1
	4	22	285	51	54	0	21	24	1
	5	30	259	137	78		4	17	1
	6	31	249	139	74	0	5	19	1
	7	30	662	182	25	1	25	49	0
	8	24	348	93	79	4	3	12	1
	9	32	219	155	8	0	1	91	0
	10	19	60	21	12	2	10	74	2
	11	8	14	20		0	10	90	
	12	33	530	29	26	16	15	43	1
	13	32	467	186	4	7	3	87	0
	14	16	68	6	23		4	72	
	15	23	200	108	86	0	4	10	1
	16	33	1316	334	4	23	1	73	0
03-May-13	1	18	233	128	75	0	1	24	
	2	18	165	118	83	0	2	15	0
	3	23	110	16	32		19	46	3
	4	23	302	62	59	0	12	28	1
	5	18	540	237	93	0	1	5	0
	6	30	257	104	71	2	10	16	1
	7	19	428	181	16	0	54	30	
	8	24	286	225	93	0	0	6	0
	9	14	35	19	29	0	5	66	
	10	10	17	6	71	1	5	24	
	11	7	22	26	19	0	0	80	
	12	20	186	37	45	0	34	20	1
	13	28	327	114	4	8	4	83	0
	14	10	59	4	0	0	1	99	
	15	20	183	124	88		1	10	1
	16	31	1155	347	7	14	2	77	0

TABLE 2. Continued

Sample Date	Transect No.	No. Species	No. Individuals	Biomass g/m2	Herb.	% Total Biomass (g)			CF
						Plankt.	Omni	Carni	
14-Jun-13	1	18	214	103	60		4	36	0
	2	18	289	238	92	0	2	6	0
	3	14	181	89	71	0	0	29	
	4	22	385	70	32	0	22	45	
	5	21	342	181	89	0	1	10	0
	6	30	229	116	81	0	5	13	1
	7	28	370	97	32	0	36	31	0
	8	24	263	89	81	0	3	15	1
	9	12	106	8	3	0	19	77	
	10	10	28	8		0	31	69	
	11	6	7	8			2	98	
	12	22	409	35	15	10	48	27	0
	13	23	468	145	2	2	1	94	0
	14	9	56	4	11	0	1	88	
	15	16	167	81	84	0	6	10	0
	16	34	816	294	47	24	1	28	0
20-Sep-13	1	21	206	90	80	0	2	18	
	2	18	129	55	81		2	17	0
	3	19	132	16	28		14	49	9
	4	22	240	24	13	1	14	72	
	5	26	324	90	68	1	7	22	3
	6	31	259	126	64	2	10	23	2
	7	28	282	95	13	2	56	28	1
	8	27	184	65	80	0	3	16	1
	9	17	266	1619	0	0	0	100	0
	10	16	46	12	1	0	11	87	1
	11	15	38	9	1	1	60	37	0
	12	31	804	372	8	3	11	78	0
	13	41	432	121	36	7	3	54	0
	14	16	154	31	56		0	43	
	15	32	259	131	83	2	4	10	1
	16	36	1554	413	46	31	1	22	0
18-Dec-13	1	22	231	115	84	0	1	14	0
	2	23	261	281	93		0	6	1
	3	26	167	64	71	0	1	25	2
	4	21	251	135	63	0	14	23	0
	5	20	165	39	66		9	23	1
	6	31	281	109	76	5	7	11	1
	7	21	337	102	16	0	51	31	2
	8	24	163	37	64	0	5	30	1
	9	23	79	13	27	3	8	54	8
	10	17	73	19	18	3	7	70	1
	11	16	76	31	24	12	17	46	0
	12	35	375	27	21	3	20	55	0
	13	37	336	107	8	1	3	88	0
	14	14	64	10	41	0	1	58	
	15	23	233	163	75	19	2	4	1
	16	36	1004	296	13	16	2	69	0

TABLE 3. Results of the Kruskal-Wallis ANOVA and the Student-Neuman-Keuls (SNK) Test addressing the question , "Are there any statistically significant differences among the mean number of fish species seen per transect,the mean number of individual fish censused per transect or the mean estimated total standing crop (in g/m2) per transect for the 15 stations among the twenty-four 2007-2013 sample periods?" The Kruskal-Wallis result is given as a "p" value at the top of the entry where (p<0.05 or less for significance). The SNK Test is used to separate means that are significantly different from one another. In the body of the table are given the sample date and mean for a given parameter on that date. Letters are used to show differences with the SNK Test; letters with the same designation show means and sample dates that are related and changes in letter designation show where significant differences exist. Overlaps in letters indicate a lack of significant differences and in such cases, only the extremes may be significantly different.

1. Mean Number of Fish Species Per Transect (p>0.28, n.s.)

Date	[n]	Mean	SNK Grouping
Jul-09	15	25.5	A
Aug-10	15	24.9	A
Aug-08	15	24.7	A
Oct-10	15	24.6	A
Nov-11	15	24.5	A
Sep-13	15	24.0	A
Nov-08	15	23.9	A
Nov-12	15	23.9	A
Jul-11	15	23.5	A
Dec-13	15	23.5	A
May-10	15	23.0	A
Dec-07	15	22.8	A
Mar-10	15	22.7	A
May-08	15	22.6	A
May-12	15	22.3	A
Jul-12	15	22.2	A
Jun-11	15	22.0	A
Mar-09	15	20.9	A
May-09	15	20.9	A
Apr-08	15	20.3	A
bMay-12	15	19.0	A
May-13	15	18.8	A
Jun-13	15	18.2	A
Feb-11	15	17.7	A

Interpretation: There are no significant differences among the mean number of species found per transect over the twenty-four sample periods.

TABLE 3. Continued.

2. Mean Number of Individual Fish Per Transect ($p>0.72$, n.s.)

YEAR	[n]	Mean	SNK Grouping
Jul-09	15	355	A
Nov-11	15	313	A
Jun-11	15	275	A
Jul-11	15	274	A
Aug-08	15	270	A
Nov-08	15	265	A
Nov-12	15	262	A
Oct-10	15	260	A
Apr-08	15	255	A
Sep-13	15	250	A
Aug-10	15	246	A
Dec-07	15	246	A
Jun-13	15	234	A
May-10	15	227	A
May-09	15	224	A
May-12	15	190	A
Jul-12	15	186	A
May-08	15	222	A
Mar-10	15	213	A
May-13	15	210	A
Dec-13	15	206	A
Mar-09	15	195	A
Feb-11	15	177	A
bMay-12	15	171	A

Interpretation: There are no significant differences among the mean number of species found per transect over the twenty-four sample periods.

3. Mean Standing Crop of Fish in g/m² Per Transect ($p>0.85$, n.s.)

YEAR	[n]	Mean	SNK Grouping
Sep-13	12	37989	A
Dec-07	12	25969	A
Nov-11	15	24047	A
Oct-10	15	23517	A
Jul-11	15	19978	A
Jul-09	15	19389	A
May-13	15	18413	A
Nov-12	15	18110	A
Jun-13	15	16859	A
Jun-11	15	16764	A
May-10	15	16578	A
Aug-10	15	16463	A
Dec-13	15	16348	A
Apr-08	15	15984	A
Nov-08	15	14174	A
May-09	15	13782	A
May-12	15	13637	A
Mar-10	15	11747	A
Mar-09	15	11705	A
Jul-12	15	11490	A
Aug-08	15	10883	A
May-08	15	10531	A
Feb-11	15	9469	A
bMay-12	15	9275	A

Interpretation: Despite the range in the estimated total standing crop per station over the twenty-four sample dates, there are no significant differences.

TABLE 4. Percent contribution based on estimated biomass for each of five feeding guilds of fishes as determined across all fifteen natural substratum stations over twenty-four survey dates in Part A. In Part B is given the same information for station 16 (Kahe outfall pipe) which was sampled commencing with the 30 May 2008 survey. In the body of the table are given the percent contribution by weight to each trophic category. Note that the December 2007 survey did not sample three of the fifteen stations. Data summarized from Table 2.

PART A. Stations 1 - 15:

Mean Percent by Weight						
Date	[n]	Herbivore	Planktivore	Omnivore	Coral Feeder	Carnivore
27-Dec-07	12	35	2	8	5	50
04-Apr-08	15	35	2	8	4	51
30-May-08	15	37	7	14	3	39
19-Aug-08	15	36	2	10	3	49
25-Nov-08	15	43	5	6	4	42
19-Mar-09	15	35	1	7	2	55
11-May-09	15	44	1	8	1	46
21-Jul-09	15	33	6	10	2	49
29-Mar-10	15	44	5	9	2	40
19-May-10	15	51	6	9	1	33
12-Aug-10	15	49	7	10	1	33
29-Oct-10	15	41	1	9	1	48
25-Feb-11	15	30	6	15	1	54
16-Jun-11	15	46	6	7	1	41
29-Jul-11	15	36	4	11	1	50
23-Nov-11	15	55	2	9	1	33
02-May-12	15	42	2	8	1	48
23-May-12	15	48	1	8	1	42
23-Jul-12	15	36	2	12	1	50
02-Nov-12	15	46	3	8	1	47
03-May-13	15	52	1	10	1	37
14-Jun-13	15	50	1	12	>1	43
20-Sep-13	15	41	2	13	2	44
18-Dec-13	15	50	4	10	1	36
<hr/>						
Grand Means		42.3	3.3	9.6	1.8	44.2

PART B. Stations 16 (Outfall Pipe) Only:

Mean Percent by Weight						
Date	[n]	Herbivore	Planktivore	Omnivore	Coral Feeder	Carnivore
30-May-08	1	8	43	2		47
19-Aug-08	1	3	55	1		41
25-Nov-08	1	10	49	1	>1	39
19-Mar-09	1	32	34	2	>1	32
11-May-09	1	35	22	6		37
21-Jul-09	1	5	36	3		56
29-Mar-10	1	24	27	2		47
19-May-10	1	13	63	8		16
12-Aug-10	1	14	27	1		58
29-Oct-10	1	7	16	1		76
25-Feb-11	1	8	32	2	>1	59
16-Jun-11	1	17	59	6	>1	17
29-Jul-11	1	4	25	1	>1	70
23-Nov-11	1	10	19	>1	>1	70
02-May-12	1	20	40	6	>1	33
23-May-12	1	14	27	2	>1	58
23-Jul-12	1	28	44	4	>1	24
02-Nov-12	1	4	23	1	>1	73
03-May-13	1	7	15	2	>1	77
14-Jun-13	1	47	24	1	>1	28
20-Sep-13	1	46	31	>1	>1	22
18-Dec-13	1	13	16	2	>1	69
<hr/>						
Grand Means		16.8	33.1	2.7	0.2	47.7

TABLE 5. Results of the Kruskal-Wallis ANOVA and the Student-Neuman-Keuls (SNK) Test addressing the question, “Are there any statistically significant differences among the mean number of fish species per transect, the mean number of individual fish per transect or the mean estimated standing crop (in g/m²) per transect among the four geographic groups of stations established on natural substratum stations 1 - 15) and sampled in the 2007-2013 period?” The four groups of transects are CIP (station nos. 1-4), Ko'Oolina (station nos. 5-7), Kahe (station nos. 8-12) and North (station nos. 13-15). The Kruskal-Wallis result is given as a “p” value at the top of the entry where (p<0.05 or less for significance). The SNK Test is used to separate means that are significantly different from one another. In the body of the table are given the four geographically-related groups of stations and parameter means per transect for each of those groups. Letters are used to show differences with the SNK Test; letters with the same designation show means and station groups that are related and changes in letter designation show where significant differences exist. Overlaps in the letters indicate a lack of significant differences and in such cases, only the extremes may be significantly different.

1. Mean Number of Fish Species Per Transect by Station Group (p<0.0001, Significant)

Station Group	(n)	Mean	SNK
			Grouping
Ko'Oolina	69	29.0	A
Nanakuli	72	21.7	B
Kahe	120	20.6	B
CIP	96	20.3	B

Interpretation: The mean number of fish species per transect at Ko'Oolina stations is significantly greater than at any of the other station groups which are all related over the twenty-four sample periods.

2. Mean Number of Individual Fish Per Transect by Station Group (p<0.0001, Significant)

Station Group	(n)	Mean	SNK
			Grouping
Ko'Oolina	69	321	A
Nanakuli	72	240	B
Kahe	120	213	B
CIP	96	210	B

Interpretation: The mean number of individual fish per transect at Ko'Oolina is significantly greater than at any of the other station groups which are all related over the twenty-four sample periods.

TABLE 5. Continued.

3. Mean Standing Crop of Fishes (in g/m²) Per Transect by Station Group (p<0.0001, Significant)

Station Group	(n)	Mean	SNK Grouping
Ko'Olina	69	131	A
CIP	96	81	B
Nanakuli	72	80	B
Kahe	120	60	B

Interpretation: Both the Kruskal-Wallis ANOVA and the SNK Test found significant differences among station groups, where the mean estimated fish standing crop was significantly greater at stations offshore of Ko'Olina than at any of the other three station groups which were all statistically related.

TABLE 6. Results of the Kruskal-Wallis ANOVA and the Student-Neuman-Keuls (SNK) Test addressing the question, "Are there any statistically significant differences among the mean number of fish species per transect, the mean number of individual fish per transect or the mean estimated standing crop (in g/m²) per transect seen among the sixteen stations established and sampled over the twenty-four periods in 2007-2013?" The Kruskal-Wallis result is given as a "p" value at the top of the entry (where p<0.05 or less for significance). The SNK Test is used to separate means that are significantly different from one another. In the body of the table are given the stations, the number of times each was sampled (n) and parameter means per transect for each. Letters are used to show differences with the SNK Test; letters with the same designation show means and station groups that are related and changes in letter designation show where significant differences exist. Overlaps in the letters indicate a lack of significant differences and in such cases, only the extremes may be significantly different.

1. Mean Number of Fish Species Per Station in 2007-13 (p < 0.0001, Significant)

Station Group	[n]	Mean	SNK Grouping			
16 (Pipe)	22	34	A			
6 (Ko'Oolina 2)	23	32	A	B		
13 (HECO 10C)	24	30		B	C	
5 (Ko'Oolina 1)	23	29		B	C	D
12 (HECO 7E)	24	28	E		C	D
8 (HECO 5B)	24	27	E	F		D
7 (HECO 1D)	23	26	E	F		
3 (East 3)	24	24		F	G	
4 (East 4)	24	23		F	G	
15 (Nana-2)	24	21			G	H
9 (HECO 7B)	24	21			G	H
2 (East 2)	24	19	I			H
10 (HECO 7C)	24	17	I			
1 (East 1)	24	16	I	J		
14 (Nana-1)	24	14		J		
11 (HECO 7D)	24	11			K	

Interpretation:

Despite the Kruskal-Wallis ANOVA finding significant differences in the mean number of fish species per transect across the sixteen stations, these differences were obscured due to overlap in the SNK Test results except for the least diverse station (number 11) which is probably related to its short length (10.5m versus 50m).

TABLE 6. Continued.

2. Mean Number of Individual Fish Censused Per Station in 2007-13 ($p < 0.0001$, Significant)

Station Group	[n]	Mean	Grouping			
16 (Pipe)	22	1162	A			
13 (HECO 10C)	24	450		B		
12 (HECO 7E)	24	398		B	C	
7 (HECO 1D)	23	360		B	C	D
8 (HECO 5B)	24	359		B	C	D
5 (Ko'Oolina 1)	23	321			C	D
4 (East 4)	24	304			C	D
6 (Ko'Oolina 2)	23	280	E			D
2 (East 2)	24	212	E	F		
3 (East 3)	24	182		F	G	
15 (Nana-2)	24	174		F	G	
9 (HECO 7B)	24	166		F	G	
1 (East 1)	24	142		F	G	
10 (HECO 7C)	24	106		F	G	H
14 (Nana-1)	24	94			G	H
11 (HECO 7D)	24	37				H

Interpretation:

The Kruskal-Wallis ANOVA noted statistically significant differences in the mean number of individual fish censused among the 16 transects over the twenty-four surveys in 2007-13. However, the SNK Test found only one clearly-obvious statistically significant station (i.e., without overlap); this was with station 16 (Kahe discharge pipe) having significantly more individual fishes present than any other and station otherwise overlap obscures further separation.

3. Mean Estimated Fish Standing Crop (g/m²) by Station in 2007-13 ($p < 0.0001$, Significant)

Station Group	[n]	Mean	Grouping			
16 (Pipe)	22	390	A			
9 (HECO 7B)	24	166		B		
6 (Ko'Oolina 2)	24	161		B	C	
13 (HECO 10C)	23	134		B	C	D
5 (Ko'Oolina 1)	23	128	E	B	C	D
7 (HECO 1D)	24	105	E	BF	C	D
15 (Nana-2)	24	97	E	BF	C	D
2 (East 2)	24	91	E	BF	C	D
3 (East 3)	24	87	E	BF	C	D
4 (East 4)	24	84	E	BF	C	D
1 (East 1)	24	61	E	F	C	D
8 (HECO 5B)	24	59	E	F	C	D
12 (HECO 7E)	24	45	E	F		D
10 (HECO 7C)	24	26	E	F		
11 (HECO 7D)	24	26	E	F		
14 (Nana-1)	24	9		F		

Interpretation:

Only one station (Kahe Discharge) had a statistically greater estimated standing crop of fishes present than found at any of the other fifteen stations whose estimated standing crops are all statistically related due to overlap.

TABLE 7. Results of the Kruskal-Wallis ANOVA and the Student-Neuman-Keuls (SNK) Test addressing the question , "Are there any statistically significant differences among the annual mean number of fish species seen per transect or the annual mean number of individual fish censused per transect among seven stations sampled in common over fifteen years encompassing a 37-year period (i.e., 1976-1984 and 2007-2013 sample periods)?" The Kruskal-Wallis result is given as a "p" value at the top of the entry where ($p < 0.05$ or less for significance). The SNK Test is used to separate means that are significantly different from one another. In the body of the table are given the sample date and mean for a given parameter on that date. Letters are used to show differences with the SNK Test; letters with the same designation show means and sample dates that are related and changes in letter designation show where significant differences exist. Overlaps in the letters indicate a lack of significant differences and in such cases, only the extremes may be significantly different.

1. Mean Number of Fish Species Per Transect ($p > 0.62$, n.s.)

YEAR	[n]	Mean	SNK Grouping
1976	3	29.0	A
1977	3	26.0	A
1979	6	24.3	A
1978	3	24.0	A
2007	7	23.7	A
2008	7	23.5	A
1984	7	23.4	A
1980	6	23.2	A
2010	7	22.3	A
2011	7	22.1	A
2012	7	21.1	A
2009	7	21.1	A
2013	7	20.6	A
1981	6	19.2	A
1982	6	17.7	A
1983	6	15.8	A

Interpretation:

There are no significant differences among the mean number of species found per transect at these seven stations among the fifteen years of sampling. Note that the highest annual means occur before the January 1980 storm event and the lowest follow that period as well as after the November 1982 hurricane.

TABLE 7. Continued.**2. Mean Number of Individual Fish Per Transect ($p>0.68$, n.s.)**

YEAR	[n]	Mean	SNK Grouping
2008	7	303.1	A
2007	7	302.3	A
2011	7	276.7	A
2009	7	271.7	A
1980	6	250.3	A
2010	7	232.3	A
2013	7	224.7	A
2012	7	215.4	A
1976	3	201.7	A
1979	6	195.0	A
1981	6	173.2	A
1978	3	169.0	A
1977	3	163.0	A
1984	7	150.0	A
1982	6	141.0	A
1983	6	85.8	A

Interpretation:

There are no significant differences among the mean number of individual fish censused per transect at these seven stations among the sixteen years of sampling. Note that the hierarchy of annual mean number of individual fish censused does not parallel that for the annual mean number of species counted at these stations. In other words, the impact of the two storm events (1980 and 1982) produced a different result with respect to the number of individual fish and the number of species counted.

FIGURE 1. Map showing the southwest coastline of Oahu from the Barbers Point Harbor on the southeast to Nanakuli Beach Park 7.9 km to the northwest. The approximate locations of each of the sixteen permanently marked stations monitored in this study are numbered. All stations except station 16 have an orientation that parallels the coastline and all are 50 m in length except for station 11 (or 7-D) which is 10.5 m in length. Station 16 is established on the terminus of the Kahe Generating facility ocean warm-water outfall and thus has an orientation that is perpendicular to the shoreline. Map courtesy of the Hawaiian Electric, Environmental Department.



Map created with TOPOGIS ©2006 National Geographic



0.0 0.5 1.0 miles
0.0 0.5 1.0 1.5 km

TN 10°
05/13/09

APPENDIX 1. Results of fish censuses carried out on each of four 2013 surveys carried out on 3 May, 14 June, 20 September and 18 December 2013. Data from the earlier surveys that comprise the first annual report are given in Brock (2009), second annual report (Brock 2010), third annual report (Brock 2011) fourth annual report (Brock 2012) and fifth annual report (Brock 2013). In the body of the table are given the list of fish species observed at each station, the trophic or feeding guild category for each species (where C=carnivore, H=herbivore, O=omnivore, P=planktivore and CF=coral feeder), the station number (here 1 through 16) as well as station name, the number of individuals of each species censused as well as the biomass (in grams) for each. Also given for each of the five trophic categories is a summary of the total number of individual fishes, the total standing crop and the percent of the total standing crop for each trophic category. Note that the total standing crop is given in grams and the area censused at each station is 200 m² except for station 11 (old Hawaiian Electric Station 7-D) which the census area is 10.5 m long and 4 m wide or 42 m². Biomass estimates for each species are based on species-specific regression coefficients using linear regression techniques (Ricker 1975, Brock and Norris 1989).

3 MAY 2013 FIELD DATA

03-May-13						GROUP	GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	1	EAST1	8	1243.38			
C	Parupeneus multifasciatus	1	EAST1	1	472.94			
C	Parupeneus multifasciatus	1	EAST1	6	576.49			
C	Plectroglyphidodon johnstonianus	1	EAST1	1	1.72			
C	Plectroglyphidodon imparipennis	1	EAST1	1	0.86			
C	Paracirrhites arcatus	1	EAST1	5	40.59			
C	Bodianus bilunulatus	1	EAST1	1	133.12			
C	Thalassoma duperrey	1	EAST1	21	575.20			
C	Thalassoma duperrey	1	EAST1	4	388.21			
C	Thalassoma duperrey	1	EAST1	9	494.54			
C	Thalassoma duperrey	1	EAST1	30	334.90			
C	Coris gaimard	1	EAST1	3	137.97			
C	Coris gaimard	1	EAST1	2	431.77			
C	Stethojulis balteata	1	EAST1	9	129.68			
C	Stethojulis balteata	1	EAST1	8	286.11			
C	Rhinecanthus rectangulus	1	EAST1	2	171.75			
C	Sufflamen fraenatus	1	EAST1	1	623.45	112	6042.67	23.6
H	Acanthurus triostegus	1	EAST1	9	416.84			
H	Acanthurus triostegus	1	EAST1	6	600.09			
H	Acanthurus nigrofuscus	1	EAST1	5	271.26			
H	Acanthurus olivaceus	1	EAST1	17	2779.78			
H	Acanthurus olivaceus	1	EAST1	18	7077.78			
H	Acanthurus olivaceus	1	EAST1	38	3563.22			
H	Acanthurus dussumieri	1	EAST1	4	2257.42			
H	Acanthurus blochii	1	EAST1	5	2267.97			
H	Naso lituratus	1	EAST1	1	14.90			
H	Naso unicornis	1	EAST1	1	38.92	104	19288.17	75.4
O	Melichthys vidua	1	EAST1	1	199.03			
O	Canthigaster jactator	1	EAST1	1	3.56	2	202.59	0.8
P	Chaetodon miliaris	1	EAST1	2	42.34			
P	Chromis vanderbilti	1	EAST1	13	4.11	15	46.44	0.2
	TOTAL	1	EAST1	233	25579.87	233	25579.87	100
C	Parupeneus multifasciatus	2	EAST2	1	155.42			
C	Plectroglyphidodon imparipennis	2	EAST2	1	0.86			
C	Paracirrhites arcatus	2	EAST2	3	24.35			
C	Bodianus bilunulatus	2	EAST2	1	1141.37			
C	Thalassoma duperrey	2	EAST2	5	485.26			
C	Thalassoma duperrey	2	EAST2	19	520.42			
C	Thalassoma duperrey	2	EAST2	11	604.44			
C	Coris venusta	2	EAST2	1	23.64			
C	Stethojulis balteata	2	EAST2	1	35.76			
C	Halichoeres ornatissimus	2	EAST2	2	32.90			
C	Rhinecanthus rectangulus	2	EAST2	3	257.62			
C	Sufflamen fraenatus	2	EAST2	1	329.34	49	3611.39	15.3
CF	Chaetodon quadrimaculatus	2	EAST2	1	62.81	1	62.81	0.3
H	Acanthurus triostegus	2	EAST2	5	500.07			
H	Acanthurus nigrofuscus	2	EAST2	2	108.50			
H	Acanthurus nigrofuscus	2	EAST2	23	326.89			
H	Acanthurus nigrofuscus	2	EAST2	22	525.73			
H	Acanthurus olivaceus	2	EAST2	8	2093.34			
H	Acanthurus olivaceus	2	EAST2	27	15205.60			
H	Naso unicornis	2	EAST2	1	525.26			
H	Naso unicornis	2	EAST2	1	278.71	89	19564.10	82.7
O	Melichthys vidua	2	EAST2	1	418.08			
O	Canthigaster jactator	2	EAST2	2	7.12	3	425.21	1.8
P	Chromis vanderbilti	2	EAST2	23	7.27	23	7.27	0.03
	TOTAL	2	EAST2	165	23670.77	165	23670.77	100

03-May-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	3	EAST3	1	54.40			
C	Parupeneus multifasciatus	3	EAST3	1	96.08			
C	Plectroglyphidodon johnstonianus	3	EAST3	6	10.33			
C	Paracirrhites arcatus	3	EAST3	5	40.59			
C	Paracirrhites forsteri	3	EAST3	1	39.65			
C	Paracirrhites forsteri	3	EAST3	1	16.35			
C	Labroides phthiropagus	3	EAST3	1	0.63			
C	Thalassoma duperrey	3	EAST3	3	291.16			
C	Thalassoma duperrey	3	EAST3	7	191.73			
C	Thalassoma duperrey	3	EAST3	6	329.69			
C	Gomphosus varius	3	EAST3	1	11.04			
C	Gomphosus varius	3	EAST3	1	22.60			
C	Coris gaimard	3	EAST3	1	83.79			
C	Stethojulis balteata	3	EAST3	4	143.05			
C	Macropharyngodon geoffroy	3	EAST3	1	5.75			
C	Halichoeres ornatissimus	3	EAST3	1	9.52			
C	Halichoeres ornatissimus	3	EAST3	1	16.45			
C	Sufflamen bursa	3	EAST3	1	85.87	43	1448.68	45.9
CF	Chaetodon quadrimaculatus	3	EAST3	1	25.30			
CF	Chaetodon multicinctus	3	EAST3	5	65.15	6	90.45	2.9
H	Calotomus carolinus	3	EAST3	2	26.93			
H	Calotomus carolinus	3	EAST3	1	72.28			
H	Scarus perspicillatus	3	EAST3	5	93.17			
H	Scarus rubroviolaceus	3	EAST3	1	85.39			
H	Acanthurus nigrofusus	3	EAST3	14	334.56			
H	Acanthurus nigrofusus	3	EAST3	18	255.83			
H	Ctenochaetus strigosus	3	EAST3	5	75.99			
H	Ctenochaetus strigosus	3	EAST3	3	80.53	49	1024.69	32.5
O	Stegastes fasciolatus	3	EAST3	3	44.29			
O	Melichthys niger	3	EAST3	2	326.67			
O	Melichthys vidua	3	EAST3	1	199.03			
O	Canthigaster jactator	3	EAST3	6	21.37	12	591.36	18.7
	TOTAL	3	EAST3	110	3155.19	110	3155.19	100
C	Parupeneus multifasciatus	4	EAST4	1	11.05			
C	Parupeneus multifasciatus	4	EAST4	1	155.42			
C	Plectroglyphidodon johnstonianus	4	EAST4	1	1.72			
C	Paracirrhites arcatus	4	EAST4	1	8.12			
C	Bodianus bilunulatus	4	EAST4	1	75.64			
C	Labroides phthiropagus	4	EAST4	2	1.26			
C	Thalassoma duperrey	4	EAST4	19	520.42			
C	Thalassoma duperrey	4	EAST4	16	879.18			
C	Thalassoma duperrey	4	EAST4	7	679.36			
C	Coris venusta	4	EAST4	1	23.64			
C	Stethojulis balteata	4	EAST4	2	71.53			
C	Macropharyngodon geoffroy	4	EAST4	8	87.83			
C	Macropharyngodon geoffroy	4	EAST4	1	81.95			
C	Halichoeres ornatissimus	4	EAST4	7	115.15			
C	Halichoeres ornatissimus	4	EAST4	1	25.14			
C	Halichoeres ornatissimus	4	EAST4	3	28.57			
C	Plagiotremus ewaensis	4	EAST4	1	0.95			
C	Zanclus cornutus	4	EAST4	3	312.47			
C	Rhinecanthus rectangulus	4	EAST4	1	85.87			
C	Sufflamen fraenatus	4	EAST4	1	224.79	78	3390.08	27.5
CF	Chaetodon ornatissimus	4	EAST4	2	138.00	2	138.00	1.1
H	Acanthurus nigrofusus	4	EAST4	13	310.66			
H	Acanthurus nigrofusus	4	EAST4	26	369.53			
H	Acanthurus olivaceus	4	EAST4	10	5631.70			
H	Naso lituratus	4	EAST4	2	897.88	51	7209.78	58.6
O	Melichthys niger	4	EAST4	5	816.68			
O	Melichthys vidua	4	EAST4	1	199.03			
O	Melichthys vidua	4	EAST4	1	418.08			
O	Cantherhines sandwichiensis	4	EAST4	1	82.05			
O	Canthigaster jactator	4	EAST4	1	3.56	9	1519.40	12.3
P	Chromis vanderbilti	4	EAST4	162	51.17	162	51.17	0.4
	TOTAL	4	EAST4	302	12308.44	302	12308.44	100

03-May-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus bifasciatus	5	KO1	1	270.20			
C	Parupeneus bifasciatus	5	KO1	1	59.77			
C	Chaetodon lunula	5	KO1	2	71.99			
C	Plectroglyphidodon johnstonianus	5	KO1	3	5.17			
C	Thalassoma duperrey	5	KO1	11	604.44			
C	Thalassoma duperrey	5	KO1	7	679.36			
C	Thalassoma duperrey	5	KO1	9	246.51			
C	Stethojulis balteata	5	KO1	1	35.76			
C	Sufflamen bursa	5	KO1	1	85.87			
C	Arothron hispidus	5	KO1	1	479.52	37	2538.59	5.3
CF	Chaetodon multicinctus	5	KO1	6	78.18	6	78.18	0.2
H	Scarus psittacus	5	KO1	1	79.27			
H	Scarus psittacus	5	KO1	1	786.15			
H	Acanthurus triostegus	5	KO1	80	8001.14			
H	Acanthurus triostegus	5	KO1	200	9263.01			
H	Acanthurus leucopareius	5	KO1	12	1592.40			
H	Acanthurus nigrofuscus	5	KO1	14	759.53			
H	Acanthurus nigrofuscus	5	KO1	24	573.53			
H	Acanthurus olivaceus	5	KO1	18	4710.01			
H	Acanthurus olivaceus	5	KO1	36	3375.68			
H	Acanthurus olivaceus	5	KO1	24	13516.09			
H	Ctenochaetus strigosus	5	KO1	19	510.03			
H	Ctenochaetus strigosus	5	KO1	18	1185.57	447	44352.42	93.4
O	Stegastes fasciolatus	5	KO1	12	177.16			
O	Melichthys niger	5	KO1	2	326.67			
O	Canthigaster jactator	5	KO1	4	14.25	18	518.08	1.1
P	Chromis vanderbilti	5	KO1	32	10.11	32	10.11	0.02
	TOTAL	5	KO1	540	47497.38	540	47497.38	100
C	Parupeneus multifasciatus	6	KO2	1	96.08			
C	Parupeneus multifasciatus	6	KO2	2	471.50			
C	Parupeneus multifasciatus	6	KO2	2	54.23			
C	Forcipiger flavissimus	6	KO2	4	36.60			
C	Plectroglyphidodon johnstonianus	6	KO2	1	1.72			
C	Cirrhitops fasciatus	6	KO2	1	8.23			
C	Bodianus bilunulatus	6	KO2	1	324.75			
C	Bodianus bilunulatus	6	KO2	1	648.58			
C	Labroides phthiropagus	6	KO2	3	1.88			
C	Thalassoma duperrey	6	KO2	7	384.64			
C	Thalassoma duperrey	6	KO2	5	485.26			
C	Thalassoma duperrey	6	KO2	8	219.12			
C	Gomphosus varius	6	KO2	1	11.04			
C	Halichoeres ornatissimus	6	KO2	2	19.05			
C	Halichoeres ornatissimus	6	KO2	1	25.14			
C	Zanclus cornutus	6	KO2	2	208.32			
C	Sufflamen bursa	6	KO2	3	257.62	45	3253.76	15.7
CF	Chaetodon unimaculatus	6	KO2	2	50.61			
CF	Chaetodon ornatissimus	6	KO2	2	138.00			
CF	Chaetodon multicinctus	6	KO2	4	52.12	8	240.73	1.2
H	Scarus psittacus	6	KO2	1	144.94			
H	Scarus psittacus	6	KO2	1	1437.46			
H	Scarus psittacus	6	KO2	5	73.07			
H	Acanthurus triostegus	6	KO2	42	1945.23			
H	Acanthurus leucopareius	6	KO2	1	66.74			
H	Acanthurus nigrofuscus	6	KO2	14	105.34			
H	Acanthurus nigrofuscus	6	KO2	24	573.53			
H	Acanthurus nigrofuscus	6	KO2	6	325.51			
H	Acanthurus nigroris	6	KO2	3	307.42			
H	Acanthurus olivaceus	6	KO2	8	4505.36			
H	Acanthurus olivaceus	6	KO2	13	3401.67			
H	Acanthurus blochii	6	KO2	2	311.17			
H	Ctenochaetus strigosus	6	KO2	4	30.30			
H	Ctenochaetus strigosus	6	KO2	7	461.05			
H	Ctenochaetus strigosus	6	KO2	9	241.60			
H	Zebrasoma flavescens	6	KO2	4	213.12			
H	Zebrasoma flavescens	6	KO2	9	85.36			
H	Zebrasoma flavescens	6	KO2	3	78.09			
H	Zebrasoma veliferum	6	KO2	2	429.30	158	14736.28	71.2
O	Melichthys niger	6	KO2	11	1796.69			
O	Melichthys vidua	6	KO2	1	199.03			
O	Cantherhines sandwichiensis	6	KO2	1	82.05			
O	Canthigaster jactator	6	KO2	3	3.68			
O	Canthigaster jactator	6	KO2	3	10.69	19	2092.14	10.1
P	Abudefduf abdominalis	6	KO2	12	375.99			
P	Chromis vanderbilti	6	KO2	15	4.74	27	380.73	1.8
	TOTAL	6	KO2	257	20703.64	257	20703.64	100

03-May-13						GROUP		GROUP PERCENT
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	
C	Mulloides flavolineatus	7	KAHE1D	2	260.56			
C	Parupeneus multifasciatus	7	KAHE1D	1	96.08			
C	Parupeneus bifasciatus	7	KAHE1D	2	788.71			
C	Plectroglyphidodon imparipennis	7	KAHE1D	3	2.59			
C	Thalassoma duperrey	7	KAHE1D	23	629.98			
C	Thalassoma duperrey	7	KAHE1D	41	2252.90			
C	Thalassoma duperrey	7	KAHE1D	66	6405.43			
C	Thalassoma ballieui	7	KAHE1D	2	205.22			
C	Gomphosus varius	7	KAHE1D	2	45.20			
C	Rhinecanthus rectangulus	7	KAHE1D	1	45.36			
C	Rhinecanthus rectangulus	7	KAHE1D	1	85.87			
C	Sufflamen bursa	7	KAHE1D	2	171.75	146	10989.66	30.4
H	Calotomus carolinus	7	KAHE1D	1	34.69			
H	Acanthurus triostegus	7	KAHE1D	2	92.63			
H	Acanthurus nigrofuscus	7	KAHE1D	70	1672.79			
H	Acanthurus nigrofuscus	7	KAHE1D	55	2983.89			
H	Ctenochaetus strigosus	7	KAHE1D	9	592.78			
H	Ctenochaetus strigosus	7	KAHE1D	8	214.75			
H	Ctenochaetus strigosus	7	KAHE1D	6	91.19	151	5682.73	15.7
O	Stegastes fasciolatus	7	KAHE1D	4	103.92			
O	Melichthys niger	7	KAHE1D	118	19273.59			
O	Cantherhines sandwichiensis	7	KAHE1D	1	82.05			
O	Canthigaster jactator	7	KAHE1D	6	21.37			
O	Canthigaster rivulata	7	KAHE1D	1	13.65	130	19494.58	53.9
P	Abudefduf abdominalis	7	KAHE1D	1	31.33	1	31.33	0.1
	TOTAL	7	KAHE1D	428	36198.30	428	36198.30	100
C	Parupeneus multifasciatus	8	KAHE5B	2	192.16			
C	Parupeneus multifasciatus	8	KAHE5B	2	108.80			
C	Plectroglyphidodon johnstonianus	8	KAHE5B	1	1.72			
C	Paracirrhites arcatus	8	KAHE5B	4	32.47			
C	Cirrhitops fasciatus	8	KAHE5B	1	8.23			
C	Thalassoma duperrey	8	KAHE5B	10	273.90			
C	Thalassoma duperrey	8	KAHE5B	8	776.42			
C	Thalassoma duperrey	8	KAHE5B	14	769.28			
C	Gomphosus varius	8	KAHE5B	1	22.60			
C	Coris venusta	8	KAHE5B	1	23.64			
C	Stethojulis balteata	8	KAHE5B	1	35.76			
C	Halichoeres ornatissimus	8	KAHE5B	1	9.52			
C	Zanclus cornutus	8	KAHE5B	2	208.32			
C	Rhinecanthus rectangulus	8	KAHE5B	1	85.87			
C	Sufflamen bursa	8	KAHE5B	3	257.62	52	2806.32	6.2
CF	Chaetodon unimaculatus	8	KAHE5B	2	50.61			
CF	Chaetodon multicinctus	8	KAHE5B	2	13.31	4	63.92	0.1
H	Cirrhipetes variolosus	8	KAHE5B	1	13.83			
H	Acanthurus triostegus	8	KAHE5B	1	100.01			
H	Acanthurus triostegus	8	KAHE5B	4	185.26			
H	Acanthurus nigrofuscus	8	KAHE5B	12	170.55			
H	Acanthurus nigrofuscus	8	KAHE5B	4	217.01			
H	Acanthurus nigrofuscus	8	KAHE5B	15	358.46			
H	Acanthurus olivaceus	8	KAHE5B	13	617.20			
H	Acanthurus olivaceus	8	KAHE5B	42	23653.15			
H	Acanthurus olivaceus	8	KAHE5B	34	8896.68			
H	Acanthurus olivaceus	8	KAHE5B	46	7521.76			
H	Naso lituratus	8	KAHE5B	1	14.90			
H	Naso lituratus	8	KAHE5B	2	48.05			
H	Naso unicornis	8	KAHE5B	1	38.92	176	41835.78	93.0
O	Stegastes fasciolatus	8	KAHE5B	5	73.82			
O	Canthigaster jactator	8	KAHE5B	2	7.12	7	80.94	0.2
P	Chromis vanderbilti	8	KAHE5B	45	14.22			
P	Naso brevirostris	8	KAHE5B	1	38.92			
P	Naso brevirostris	8	KAHE5B	1	123.12	47	176.26	0.4
	TOTAL	8	KAHE5B	286	44963.21	286	44963.21	100
C	Mulloides flavolineatus	9	KAHE7B	1	965.72			
C	Parupeneus pleurostigma	9	KAHE7B	2	116.02			
C	Parupeneus pleurostigma	9	KAHE7B	1	153.39			
C	Parupeneus pleurostigma	9	KAHE7B	1	98.25			
C	Parupeneus multifasciatus	9	KAHE7B	1	155.42			
C	Thalassoma duperrey	9	KAHE7B	1	27.39			
C	Thalassoma duperrey	9	KAHE7B	4	388.21			
C	Thalassoma duperrey	9	KAHE7B	2	109.90			
C	Coris venusta	9	KAHE7B	1	23.64			
C	Coris venusta	9	KAHE7B	1	48.38			
C	Zanclus cornutus	9	KAHE7B	1	104.16			
C	Sufflamen bursa	9	KAHE7B	1	85.87			
C	Sufflamen bursa	9	KAHE7B	2	289.30	19	2565.64	66.0
H	Acanthurus nigrofuscus	9	KAHE7B	1	23.90			
H	Acanthurus olivaceus	9	KAHE7B	4	654.07			
H	Naso lituratus	9	KAHE7B	1	72.58			
H	Naso unicornis	9	KAHE7B	3	369.35	9	1119.90	28.8
O	Melichthys vidua	9	KAHE7B	1	199.03			
O	Canthigaster jactator	9	KAHE7B	1	2.21	2	201.24	5.2
P	Chromis vanderbilti	9	KAHE7B	5	1.58	5	1.58	0.04
	TOTAL	9	KAHE7B	35	3888.36	35	3888.36	100

03-May-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Chromis vanderbilti	10	KAHE7C	3	0.95			
C	Thalassoma duperrey	10	KAHE7C	1	3.15			
C	Coris venusta	10	KAHE7C	1	48.38			
C	Acanthurus triostegus	10	KAHE7C	1	100.01			
C	Sufflamen bursa	10	KAHE7C	1	144.65	7	297.14	23.8
H	Coris gaimard	10	KAHE7C	1	139.13			
H	Acanthurus olivaceus	10	KAHE7C	3	490.55			
H	Sufflamen bursa	10	KAHE7C	3	257.62	7	887.30	71.1
O	Acanthurus blochii	10	KAHE7C	1	56.70	1	56.70	4.5
P	Canthigaster jactator	10	KAHE7C	2	7.12	2	7.12	0.6
	TOTAL	10	KAHE7C	17	1248.27	17	1248.27	100
C	Paracirrhites arcatus	11	KAHE7D	2	16.24			
C	Coris gaimard	11	KAHE7D	2	636.12			
C	Sufflamen bursa	11	KAHE7D	1	85.87			
C	Sufflamen bursa	11	KAHE7D	1	144.65	6	882.88	80.2
H	Acanthurus triostegus	11	KAHE7D	1	46.32			
H	Acanthurus olivaceus	11	KAHE7D	1	163.52	2	209.83	19.1
O	Canthigaster jactator	11	KAHE7D	1	3.56	1	3.56	0.3
P	Chromis vanderbilti	11	KAHE7D	13	4.11	13	4.11	0.4
	TOTAL	11	KAHE7D	22	1100.38	22	1100.38	100
C	Parupeneus multifasciatus	12	KAHE7E	2	192.16			
C	Parupeneus multifasciatus	12	KAHE7E	3	163.20			
C	Parupeneus multifasciatus	12	KAHE7E	2	54.23			
C	Plectroglyphidodon johnstonianus	12	KAHE7E	2	3.44			
C	Paracirrhites arcatus	12	KAHE7E	2	16.24			
C	Thalassoma duperrey	12	KAHE7E	2	109.90			
C	Thalassoma duperrey	12	KAHE7E	1	97.05			
C	Pseudojuloides cerasinus	12	KAHE7E	2	54.78			
C	Pseudojuloides cerasinus	12	KAHE7E	7	22.05			
C	Zanclus cornutus	12	KAHE7E	1	104.16			
C	Sufflamen bursa	12	KAHE7E	4	343.49			
C	Sufflamen fraenatus	12	KAHE7E	1	329.34	29	1490.05	20.3
CF	Chaetodon multicinctus	12	KAHE7E	6	39.94	6	39.94	0.5
H	Calotomus carolinus	12	KAHE7E	1	34.69			
H	Acanthurus triostegus	12	KAHE7E	1	100.01			
H	Acanthurus nigrofusus	12	KAHE7E	5	71.06			
H	Acanthurus nigrofusus	12	KAHE7E	9	67.72			
H	Acanthurus olivaceus	12	KAHE7E	3	1689.51			
H	Acanthurus olivaceus	12	KAHE7E	3	281.31			
H	Acanthurus olivaceus	12	KAHE7E	1	0.69			
H	Acanthurus olivaceus	12	KAHE7E	4	1046.67			
H	Naso unicornis	12	KAHE7E	1	17.19	28	3308.86	45.1
O	Melichthys niger	12	KAHE7E	6	980.01			
O	Melichthys vidua	12	KAHE7E	7	1393.21			
O	Cantherhines sandwichiensis	12	KAHE7E	1	82.05			
O	Canthigaster jactator	12	KAHE7E	4	8.82	18	2464.10	33.6
P	Chromis vanderbilti	12	KAHE7E	102	32.22			
P	Chromis hanui	12	KAHE7E	3	2.24	105	34.46	0.5
	TOTAL	12	KAHE7E	186	7337.40	186	7337.40	100
C	Saurida gracilis	13	KAHE10	2	366.52			
C	Aulostomus chinensis	13	KAHE10	1	27.72			
C	Lutjanus kasmira	13	KAHE10	72	13726.57			
C	Lutjanus kasmira	13	KAHE10	40	2294.10			
C	Parupeneus multifasciatus	13	KAHE10	1	155.42			
C	Parupeneus bifasciatus	13	KAHE10	1	59.77			
C	Forcipiger flavissimus	13	KAHE10	1	9.15			
C	Paracirrhites arcatus	13	KAHE10	2	16.24			
C	Thalassoma duperrey	13	KAHE10	10	970.52			
C	Thalassoma duperrey	13	KAHE10	7	22.05			
C	Thalassoma duperrey	13	KAHE10	8	219.12			
C	Thalassoma duperrey	13	KAHE10	12	659.39			
C	Thalassoma ballieui	13	KAHE10	1	102.61			
C	Coris venusta	13	KAHE10	1	86.86			
C	Coris gaimard	13	KAHE10	2	91.98			
C	Stethojulis balteata	13	KAHE10	1	35.76			
C	Macropharyngodon geoffroy	13	KAHE10	1	18.63			
C	Sufflamen bursa	13	KAHE10	1	85.87	164	18948.30	83.0
CF	Chaetodon multicinctus	13	KAHE10	2	26.06	2	26.06	0.1
H	Acanthurus triostegus	13	KAHE10	1	46.32			
H	Acanthurus leucopareius	13	KAHE10	1	27.52			
H	Acanthurus nigrofusus	13	KAHE10	5	119.49			
H	Acanthurus nigrofusus	13	KAHE10	8	434.02			
H	Acanthurus olivaceus	13	KAHE10	1	163.52			
H	Acanthurus olivaceus	13	KAHE10	3	142.43			
H	Ctenochaetus strigosus	13	KAHE10	1	15.20	20	948.48	4.2
O	Stegastes fasciolatus	13	KAHE10	25	369.08			
O	Melichthys vidua	13	KAHE10	3	597.09			
O	Canthigaster coronata	13	KAHE10	1	13.65	29	979.82	4.3
P	Dascyllus albisella	13	KAHE10	21	64.85			
P	Abudefduf abdominalis	13	KAHE10	18	563.99			
P	Abudefduf vaigensis	13	KAHE10	15	702.78			
P	Chromis ovalis	13	KAHE10	12	71.18			
P	Chromis ovalis	13	KAHE10	45	520.19			
P	Naso brevirostris	13	KAHE10	1	10.25	112	1933.23	8.5
	TOTAL	13	KAHE10	327	22835.89	327	22835.89	100

03-May-13						GROUP		GROUP PERCENT
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	
C	Parupeneus multifasciatus	14	NANA1	1	11.05			
C	Plectroglyphidodon imparipennis	14	NANA1	4	3.45			
C	Cirrhitops fasciatus	14	NANA1	1	8.23			
C	Thalassoma duperrey	14	NANA1	4	109.56			
C	Thalassoma duperrey	14	NANA1	4	219.80			
C	Thalassoma duperrey	14	NANA1	2	194.10			
C	Thalassoma duperrey	14	NANA1	10	111.63			
C	Thalassoma duperrey	14	NANA1	12	37.81			
C	Stethojulis balteata	14	NANA1	1	72.39			
C	Rhinecanthus rectangulus	14	NANA1	2	90.72			
C	Ostracion meleagris	14	NANA1	1	6.76			
C	Ostracion meleagris	14	NANA1	1	2.71	43	868.21	98.7
H	Acanthurus blochii	14	NANA1	1	0.45	1	0.45	0.1
O	Canthigaster jactator	14	NANA1	2	7.12	2	7.12	0.8
P	Chromis vanderbilti	14	NANA1	13	4.11	13	4.11	0.5
	TOTAL	14	NANA1	59	879.90	59	879.90	100
C	Adioryx spinifer	15	NANA2	1	265.90			
C	Monotaxis grandoculis	15	NANA2	2	27.47			
C	Parupeneus multifasciatus	15	NANA2	1	27.12			
C	Paracirrhites forsteri	15	NANA2	2	79.29			
C	Thalassoma duperrey	15	NANA2	11	604.44			
C	Thalassoma duperrey	15	NANA2	11	1067.57			
C	Thalassoma duperrey	15	NANA2	8	219.12			
C	Macropharyngodon geoffroy	15	NANA2	1	18.63			
C	Halichoeres ornatissimus	15	NANA2	1	16.45			
C	Rhinecanthus rectangulus	15	NANA2	1	45.36			
C	Sufflamen bursa	15	NANA2	1	85.87	40	2457.23	9.9
CF	Chaetodon ornatissimus	15	NANA2	2	273.78			
CF	Chaetodon multicinctus	15	NANA2	6	39.94	8	313.72	1.3
H	Acanthurus leucopareus	15	NANA2	9	1194.30			
H	Acanthurus nigrofusus	15	NANA2	14	334.56			
H	Acanthurus nigrofusus	15	NANA2	4	56.85			
H	Acanthurus nigrofusus	15	NANA2	5	271.26			
H	Acanthurus olivaceus	15	NANA2	25	2344.22			
H	Acanthurus olivaceus	15	NANA2	18	10137.06			
H	Acanthurus olivaceus	15	NANA2	23	6018.34			
H	Acanthurus blochii	15	NANA2	12	680.39			
H	Acanthurus blochii	15	NANA2	3	466.75			
H	Ctenochaetus strigosus	15	NANA2	5	134.22			
H	Zebrasoma flavescens	15	NANA2	2	106.56	120	21744.52	87.6
O	Stegastes fasciolatus	15	NANA2	9	132.87			
O	Melichthys niger	15	NANA2	1	163.34			
O	Canthigaster jactator	15	NANA2	5	17.81	15	314.02	1.3
	TOTAL	15	NANA2	183	24829.49	183	24829.49	100
C	Aulostomus chinensis	16	PIPE	1	88.50			
C	Decapterus macarellus	16	PIPE	121	32590.24			
C	Decapterus macarellus	16	PIPE	15	1954.19			
C	Lutjanus kasmira	16	PIPE	36	930.86			
C	Lutjanus kasmira	16	PIPE	41	2351.45			
C	Parupeneus multifasciatus	16	PIPE	1	835.32			
C	Parupeneus multifasciatus	16	PIPE	1	235.75			
C	Parupeneus multifasciatus	16	PIPE	5	480.41			
C	Forcipiger flavissimus	16	PIPE	2	18.30			
C	Plectroglyphidodon johnstonianus	16	PIPE	1	1.72			
C	Labroides phthirophagus	16	PIPE	2	1.26			
C	Thalassoma duperrey	16	PIPE	123	11937.39			
C	Thalassoma ballieui	16	PIPE	2	335.05			
C	Thalassoma ballieui	16	PIPE	1	520.80			
C	Gomphosus varius	16	PIPE	3	186.09			
C	Gomphosus varius	16	PIPE	8	180.81			
C	Stethojulis balteata	16	PIPE	3	217.17			
C	Macropharyngodon geoffroy	16	PIPE	4	74.51			
C	Halichoeres ornatissimus	16	PIPE	2	32.90			
C	Zanclus cornutus	16	PIPE	2	208.32			
C	Sufflamen bursa	16	PIPE	4	343.49	378	53524.53	77.0
CF	Chaetodon multicinctus	16	PIPE	2	13.31			
CF	Exallias brevis	16	PIPE	1	49.35	3	62.66	0.1
H	Scarus sordidus	16	PIPE	9	1260.05			
H	Scarus sordidus	16	PIPE	1	775.07			
H	Scarus sordidus	16	PIPE	1	234.65			
H	Scarus sordidus	16	PIPE	6	456.08			
H	Scarus psittacus	16	PIPE	1	79.27			
H	Acanthurus nigrofusus	16	PIPE	6	143.38			
H	Acanthurus nigrofusus	16	PIPE	5	271.26			
H	Acanthurus blochii	16	PIPE	1	56.70			
H	Ctenochaetus strigosus	16	PIPE	11	724.51			
H	Ctenochaetus strigosus	16	PIPE	12	322.13			
H	Naso unicornis	16	PIPE	1	38.92			
H	Naso unicornis	16	PIPE	2	381.49	56	4743.52	6.8
O	Stegastes fasciolatus	16	PIPE	41	605.29			
O	Melichthys vidua	16	PIPE	4	497.38			
O	Canthigaster jactator	16	PIPE	5	17.81	50	1120.49	1.6
P	Chaetodon miliaris	16	PIPE	25	312.45			
P	Abudefduf abdominalis	16	PIPE	48	1503.96			
P	Abudefduf vaigensis	16	PIPE	90	4216.67			
P	Chromis vanderbilti	16	PIPE	115	36.33			
P	Chromis ovalis	16	PIPE	95	563.52			
P	Chromis ovalis	16	PIPE	295	3410.10	668	10043.04	14.5
	TOTAL	16	PIPE	1155	69494.23	1155	69494.23	100

14 JUNE 2013 FIELD DATA

14-Jun-13

GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	GROUP BIOMASS	GROUP PERCENT
C	Parupeneus multifasciatus	1	EAST1	3	81.35			
C	Parupeneus multifasciatus	1	EAST1	4	942.99			
C	Parupeneus multifasciatus	1	EAST1	3	1021.33			
C	Parupeneus multifasciatus	1	EAST1	6	932.53			
C	Plectroglyphidodon imparipennis	1	EAST1	1	0.86			
C	Paracirrhites arcatus	1	EAST1	2	16.24			
C	Paracirrhites forsteri	1	EAST1	1	16.35			
C	Bodianus bilunulatus	1	EAST1	1	75.64			
C	Labroides phthirophagus	1	EAST1	2	1.26			
C	Thalassoma duperrey	1	EAST1	25	2426.30			
C	Thalassoma duperrey	1	EAST1	17	934.13			
C	Thalassoma duperrey	1	EAST1	16	50.41			
C	Thalassoma duperrey	1	EAST1	12	133.96			
C	Coris gaimard	1	EAST1	1	215.88			
C	Stethojulis balteata	1	EAST1	5	178.82			
C	Stethojulis balteata	1	EAST1	8	115.27			
C	Sufflamen fraenatus	1	EAST1	1	224.79	108	7368.11	35.6
CF	Chaetodon quadrimaculatus	1	EAST1	2	50.61	2	50.61	0.2
H	Acanthurus triostegus	1	EAST1	18	833.67			
H	Acanthurus nigrofusus	1	EAST1	31	740.81			
H	Acanthurus nigrofusus	1	EAST1	14	759.53			
H	Acanthurus olivaceus	1	EAST1	12	6758.04			
H	Acanthurus olivaceus	1	EAST1	9	843.92			
H	Acanthurus olivaceus	1	EAST1	3	142.43			
H	Acanthurus olivaceus	1	EAST1	5	98.72			
H	Acanthurus blochii	1	EAST1	4	928.96			
H	Naso lituratus	1	EAST1	2	1244.71	98	12350.79	59.8
O	Melichthys vidua	1	EAST1	3	888.72			
O	Canthigaster jactator	1	EAST1	3	10.69	6	899.40	4.4
	TOTAL	1	EAST1	214	20668.92	214	20668.92	100
C	Parupeneus multifasciatus	2	EAST2	1	54.40			
C	Parupeneus multifasciatus	2	EAST2	2	54.23			
C	Parupeneus multifasciatus	2	EAST2	1	235.75			
C	Plectroglyphidodon johnstonianus	2	EAST2	1	1.72			
C	Paracirrhites arcatus	2	EAST2	1	8.12			
C	Labroides phthirophagus	2	EAST2	2	1.26			
C	Thalassoma duperrey	2	EAST2	19	1044.03			
C	Thalassoma duperrey	2	EAST2	14	156.29			
C	Thalassoma duperrey	2	EAST2	14	383.47			
C	Thalassoma duperrey	2	EAST2	7	679.36			
C	Gomphosus varius	2	EAST2	3	33.12			
C	Halichoeres ornatissimus	2	EAST2	1	16.45			
C	Halichoeres ornatissimus	2	EAST2	1	25.14			
C	Rhinecanthus rectangulus	2	EAST2	2	171.75	69	2865.08	6.0
CF	Chaetodon unimaculatus	2	EAST2	2	50.61			
CF	Chaetodon quadrimaculatus	2	EAST2	1	25.30			
CF	Chaetodon quadrimaculatus	2	EAST2	1	0.79	4	76.70	0.2
H	Acanthurus nigrofusus	2	EAST2	47	1123.16			
H	Acanthurus nigrofusus	2	EAST2	32	454.81			
H	Acanthurus olivaceus	2	EAST2	40	10466.68			
H	Acanthurus olivaceus	2	EAST2	50	28158.51			
H	Acanthurus olivaceus	2	EAST2	12	1962.20			
H	Acanthurus blochii	2	EAST2	4	622.33			
H	Acanthurus blochii	2	EAST2	1	330.67			
H	Naso lituratus	2	EAST2	1	622.36	187	43740.72	91.8
O	Stegastes fasciolatus	2	EAST2	4	59.05			
O	Melichthys vidua	2	EAST2	3	888.72			
O	Canthigaster jactator	2	EAST2	3	10.69	10	958.45	2.0
P	Chromis vanderbilti	2	EAST2	19	6.00	19	6.00	0.01
	TOTAL	2	EAST2	289	47646.95	289	47646.95	100

14-Jun-13						GROUP		GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	3	EAST3	8	88.41			
C	Plectroglyphidodon johnstonianus	3	EAST3	3	5.17			
C	Paracirrhites arcatus	3	EAST3	6	48.71			
C	Thalassoma duperrey	3	EAST3	25	1373.72			
C	Thalassoma duperrey	3	EAST3	22	245.59			
C	Thalassoma duperrey	3	EAST3	16	1552.83			
C	Thalassoma duperrey	3	EAST3	12	328.69			
C	Coris gaimard	3	EAST3	1	139.13			
C	Stethojulis balteata	3	EAST3	2	144.78			
C	Halichoeres ornatissimus	3	EAST3	1	9.52			
C	Halichoeres ornatissimus	3	EAST3	1	25.14			
C	Rhinecanthus rectangulus	3	EAST3	2	90.72			
C	Rhinecanthus rectangulus	3	EAST3	2	171.75			
C	Sufflamen fraenatus	3	EAST3	1	461.25			
C	Sufflamen fraenatus	3	EAST3	1	329.34			
C	Sufflamen fraenatus	3	EAST3	1	224.79	104	5239.54	29.3
H	Acanthurus triostegus	3	EAST3	16	741.04			
H	Acanthurus triostegus	3	EAST3	16	1600.23			
H	Acanthurus olivaceus	3	EAST3	15	8447.55			
H	Acanthurus olivaceus	3	EAST3	6	1570.00			
H	Acanthurus blochii	3	EAST3	1	232.24	54	12591.07	70.5
O	Canthigaster jactator	3	EAST3	6	21.37	6	21.37	0.1
P	Chromis vanderbilti	3	EAST3	17	5.37	17	5.37	0.03
	TOTAL	3	EAST3	181	17857.35	181	17857.35	99.9999999
C	Cephalopholis argus	4	EAST4	1	638.29			
C	Parupeneus multifasciatus	4	EAST4	12	132.62			
C	Parupeneus multifasciatus	4	EAST4	2	108.80			
C	Plectroglyphidodon johnstonianus	4	EAST4	2	3.44			
C	Paracirrhites arcatus	4	EAST4	9	73.06			
C	Paracirrhites forsteri	4	EAST4	1	9.32			
C	Bodianus bilunulatus	4	EAST4	1	133.12			
C	Bodianus bilunulatus	4	EAST4	1	467.86			
C	Labroides phthirophagus	4	EAST4	2	1.26			
C	Pseudocheilinus octotaenia	4	EAST4	1	14.41			
C	Thalassoma duperrey	4	EAST4	14	1358.73			
C	Thalassoma duperrey	4	EAST4	24	1318.77			
C	Thalassoma duperrey	4	EAST4	34	931.28			
C	Thalassoma duperrey	4	EAST4	37	413.04			
C	Coris venusta	4	EAST4	5	241.90			
C	Coris venusta	4	EAST4	3	70.91			
C	Macropharyngodon geoffroy	4	EAST4	3	17.24			
C	Halichoeres ornatissimus	4	EAST4	3	49.35			
C	Halichoeres ornatissimus	4	EAST4	2	19.05			
C	Halichoeres ornatissimus	4	EAST4	2	50.27			
C	Sufflamen fraenatus	4	EAST4	1	224.79	160	6277.51	45.1
H	Acanthurus nigrofusus	4	EAST4	42	596.93			
H	Acanthurus nigrofusus	4	EAST4	28	669.12			
H	Acanthurus olivaceus	4	EAST4	5	2815.85			
H	Acanthurus dussumieri	4	EAST4	1	434.70	76	4516.60	32.5
O	Melichthys niger	4	EAST4	7	1736.22			
O	Melichthys vidua	4	EAST4	3	888.72			
O	Melichthys vidua	4	EAST4	1	199.03			
O	Cantherhines sandwichiensis	4	EAST4	1	131.57			
O	Cantherhines sandwichiensis	4	EAST4	1	82.05			
O	Canthigaster coronata	4	EAST4	1	7.59			
O	Canthigaster jactator	4	EAST4	8	28.50	22	3073.68	22.1
P	Chromis vanderbilti	4	EAST4	127	40.12	127	40.12	0.3
	TOTAL	4	EAST4	385	13907.91	385	13907.91	99.9999997

14-Jun-13						GROUP		GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	5	KO1	2	310.84			
C	Parupeneus multifasciatus	5	KO1	1	27.12			
C	Parupeneus multifasciatus	5	KO1	2	108.80			
C	Parupeneus bifasciatus	5	KO1	1	176.01			
C	Parupeneus bifasciatus	5	KO1	2	540.40			
C	Plectroglyphidodon johnstonianus	5	KO1	6	10.33			
C	Labroides phthirophagus	5	KO1	3	1.88			
C	Thalassoma duperrey	5	KO1	7	679.36			
C	Thalassoma duperrey	5	KO1	16	879.18			
C	Thalassoma duperrey	5	KO1	5	136.95			
C	Thalassoma duperrey	5	KO1	11	122.80			
C	Thalassoma ballieui	5	KO1	1	167.53			
C	Gomphosus varius	5	KO1	1	22.60			
C	Gomphosus varius	5	KO1	1	62.03			
C	Gomphosus varius	5	KO1	1	91.05			
C	Sufflamen bursa	5	KO1	1	144.65			
C	Sufflamen bursa	5	KO1	1	85.87			
C	Ostracion meleagris	5	KO1	1	6.76	63	3574.17	9.9
CF	Chaetodon multicinctus	5	KO1	4	52.12	4	52.12	0.1
H	Acanthurus triostegus	5	KO1	11	509.47			
H	Acanthurus leucopareius	5	KO1	5	1163.38			
H	Acanthurus nigrofusus	5	KO1	17	922.29			
H	Acanthurus nigrofusus	5	KO1	14	334.56			
H	Acanthurus nigrofusus	5	KO1	7	52.67			
H	Acanthurus olivaceus	5	KO1	13	1218.99			
H	Acanthurus olivaceus	5	KO1	12	6758.04			
H	Acanthurus olivaceus	5	KO1	9	427.29			
H	Acanthurus olivaceus	5	KO1	41	10728.35			
H	Acanthurus blochii	5	KO1	9	881.79			
H	Acanthurus blochii	5	KO1	7	1625.68			
H	Acanthurus blochii	5	KO1	3	87.09			
H	Ctenochaetus strigosus	5	KO1	23	3039.08			
H	Ctenochaetus strigosus	5	KO1	33	885.85			
H	Ctenochaetus strigosus	5	KO1	35	2305.27			
H	Zebrasoma flavescens	5	KO1	5	266.40			
H	Naso lituratus	5	KO1	3	617.96			
H	Naso lituratus	5	KO1	1	448.94	248	32273.11	89.4
O	Stegastes fasciatus	5	KO1	7	181.86			
O	Canthigaster jactator	5	KO1	8	28.50	15	210.36	0.6
P	Chromis vanderbilti	5	KO1	12	3.79	12	3.79	0.01
	TOTAL	5	KO1	342	36113.55	342	36113.55	100
C	Cephalopholis argus	6	KO2	1	638.29			
C	Parupeneus multifasciatus	6	KO2	2	192.16			
C	Parupeneus cyclostomus	6	KO2	1	316.39			
C	Plectroglyphidodon johnstonianus	6	KO2	1	1.72			
C	Labroides phthirophagus	6	KO2	3	1.88			
C	Thalassoma duperrey	6	KO2	2	194.10			
C	Thalassoma duperrey	6	KO2	7	78.14			
C	Thalassoma duperrey	6	KO2	7	191.73			
C	Thalassoma duperrey	6	KO2	5	274.74			
C	Gomphosus varius	6	KO2	3	33.12			
C	Stethojulis balteata	6	KO2	1	35.76			
C	Zanclus cornutus	6	KO2	2	208.32			
C	Sufflamen bursa	6	KO2	1	144.65			
C	Sufflamen bursa	6	KO2	1	45.36			
C	Sufflamen fraenatus	6	KO2	1	623.45	38	2979.84	12.8
CF	Chaetodon unimaculatus	6	KO2	2	50.61			
CF	Chaetodon ornatissimus	6	KO2	2	138.00			
CF	Chaetodon multicinctus	6	KO2	4	52.12	8	240.73	1.0
H	Scarus psittacus	6	KO2	1	14.61			
H	Scarus psittacus	6	KO2	1	375.61			
H	Scarus rubroviolaceus	6	KO2	1	2733.87			
H	Acanthurus triostegus	6	KO2	7	324.21			
H	Acanthurus triostegus	6	KO2	1	4.24			
H	Acanthurus leucopareius	6	KO2	6	796.20			
H	Acanthurus nigrofusus	6	KO2	5	37.62			
H	Acanthurus nigrofusus	6	KO2	19	454.04			
H	Acanthurus nigrofusus	6	KO2	12	170.55			
H	Acanthurus nigroris	6	KO2	3	71.69			
H	Acanthurus olivaceus	6	KO2	9	1471.65			
H	Acanthurus olivaceus	6	KO2	13	7321.21			
H	Acanthurus blochii	6	KO2	5	777.91			
H	Acanthurus blochii	6	KO2	2	24.49			
H	Ctenochaetus strigosus	6	KO2	10	75.76			
H	Ctenochaetus strigosus	6	KO2	10	658.65			
H	Ctenochaetus strigosus	6	KO2	13	348.97			
H	Ctenochaetus strigosus	6	KO2	12	1585.60			
H	Zebrasoma flavescens	6	KO2	6	156.18			
H	Zebrasoma flavescens	6	KO2	8	75.87			
H	Zebrasoma flavescens	6	KO2	16	852.48			
H	Naso lituratus	6	KO2	1	448.94	161	18780.38	81.0
O	Melichthys niger	6	KO2	2	496.06			
O	Melichthys vidua	6	KO2	1	418.08			
O	Melichthys vidua	6	KO2	1	199.03			
O	Canthigaster jactator	6	KO2	3	10.69	7	1123.86	4.8
P	Abudefduf abdominalis	6	KO2	2	62.67			
P	Chromis vanderbilti	6	KO2	13	4.11	15	66.77	0.3
	TOTAL	6	KO2	229	23191.58	229	23191.58	99.9999998

14-Jun-13						GROUP		GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Saurida gracilis	7	KAHE1D	1	83.87			
C	Myripristis amaenus	7	KAHE1D	3	246.25			
C	Fistularia commersoni	7	KAHE1D	1	165.95			
C	Parupeneus porphyreus	7	KAHE1D	1	185.13			
C	Parupeneus multifasciatus	7	KAHE1D	1	96.08			
C	Chaetodon fremblii	7	KAHE1D	1	20.91			
C	Plectroglyphidodon johnstonianus	7	KAHE1D	3	5.17			
C	Paracirrhites arcatus	7	KAHE1D	1	16.35			
C	Labroides phthirophagus	7	KAHE1D	2	1.26			
C	Thalassoma duperrey	7	KAHE1D	15	824.23			
C	Thalassoma duperrey	7	KAHE1D	25	684.76			
C	Thalassoma duperrey	7	KAHE1D	30	2911.56			
C	Thalassoma duperrey	7	KAHE1D	12	133.96			
C	Coris gaimard	7	KAHE1D	1	22.07			
C	Coris gaimard	7	KAHE1D	4	34.26			
C	Stethojulis balteata	7	KAHE1D	3	107.29			
C	Stethojulis balteata	7	KAHE1D	5	20.01			
C	Rhinecanthus rectangulus	7	KAHE1D	1	85.87			
C	Sufflamen bursa	7	KAHE1D	1	45.36			
C	Sufflamen bursa	7	KAHE1D	4	343.49	115	6033.84	31.1
CF	Chaetodon ornatissimus	7	KAHE1D	2	57.06			
CF	Chaetodon multicinctus	7	KAHE1D	2	13.31	4	70.37	0.4
H	Centropyge potteri	7	KAHE1D	1	10.62			
H	Scarus psittacus	7	KAHE1D	31	453.04			
H	Scarus psittacus	7	KAHE1D	49	1855.76			
H	Acanthurus triostegus	7	KAHE1D	9	154.50			
H	Acanthurus triostegus	7	KAHE1D	21	972.62			
H	Acanthurus nigrofuscus	7	KAHE1D	32	764.71			
H	Acanthurus nigrofuscus	7	KAHE1D	8	60.20			
H	Acanthurus olivaceus	7	KAHE1D	9	427.29			
H	Acanthurus olivaceus	7	KAHE1D	12	1125.23			
H	Ctenochaetus strigosus	7	KAHE1D	12	322.13			
H	Zebbrasoma flavescens	7	KAHE1D	2	106.56	186	6252.65	32.3
O	Stegastes fasciolatus	7	KAHE1D	10	259.80			
O	Melichthys niger	7	KAHE1D	40	6533.42			
O	Melichthys vidua	7	KAHE1D	1	199.03			
O	Canthigaster jactator	7	KAHE1D	7	24.93	58	7017.19	36.2
P	Chromis vanderbilti	7	KAHE1D	7	2.21	7	2.21	0.01
	TOTAL	7	KAHE1D	370	19376.26	370	19376.26	100
C	Parupeneus pleurostigma	8	KAHE5B	1	58.01			
C	Parupeneus multifasciatus	8	KAHE5B	15	165.77			
C	Parupeneus multifasciatus	8	KAHE5B	1	54.40			
C	Parupeneus multifasciatus	8	KAHE5B	2	310.84			
C	Plectroglyphidodon johnstonianus	8	KAHE5B	5	8.61			
C	Plectroglyphidodon imparipennis	8	KAHE5B	1	0.86			
C	Paracirrhites arcatus	8	KAHE5B	3	24.35			
C	Thalassoma duperrey	8	KAHE5B	7	679.36			
C	Thalassoma duperrey	8	KAHE5B	10	549.49			
C	Thalassoma duperrey	8	KAHE5B	8	89.31			
C	Thalassoma duperrey	8	KAHE5B	12	328.69			
C	Halichoeres ornatissimus	8	KAHE5B	1	16.45			
C	Halichoeres ornatissimus	8	KAHE5B	1	9.52			
C	Zanclus cornutus	8	KAHE5B	1	104.16			
C	Rhinecanthus rectangulus	8	KAHE5B	2	171.75			
C	Sufflamen bursa	8	KAHE5B	2	171.75	72	2743.32	15.5
CF	Chaetodon multicinctus	8	KAHE5B	2	26.06			
CF	Exallias brevis	8	KAHE5B	1	49.35			
CF	Pervagor melanocephalus	8	KAHE5B	1	16.22	4	91.62	0.5
H	Calotomus carolinus	8	KAHE5B	1	34.69			
H	Acanthurus nigrofuscus	8	KAHE5B	18	430.15			
H	Acanthurus nigrofuscus	8	KAHE5B	14	198.98			
H	Acanthurus nigrofuscus	8	KAHE5B	20	150.49			
H	Acanthurus olivaceus	8	KAHE5B	13	1218.99			
H	Acanthurus olivaceus	8	KAHE5B	19	10700.23			
H	Acanthurus olivaceus	8	KAHE5B	6	1570.00			
H	Acanthurus blochii	8	KAHE5B	1	12.25			
H	Naso lituratus	8	KAHE5B	1	72.58			
H	Naso unicornis	8	KAHE5B	1	17.19	94	14405.57	81.1
O	Stegastes fasciolatus	8	KAHE5B	1	25.98			
O	Melichthys vidua	8	KAHE5B	1	199.03			
O	Cantherhines sandwichiensis	8	KAHE5B	3	246.15			
O	Canthigaster jactator	8	KAHE5B	5	17.81	10	488.97	2.8
P	Chromis vanderbilti	8	KAHE5B	83	26.22	83	26.22	0.1
	TOTAL	8	KAHE5B	263	17755.70	263	17755.70	100
C	Gymnothorax meleagris	9	KAHE7B	1	57.93			
C	Saurida gracilis	9	KAHE7B	1	51.93			
C	Lutjanus kasmira	9	KAHE7B	70	54.57			
C	Parupeneus pleurostigma	9	KAHE7B	8	464.07			
C	Parupeneus multifasciatus	9	KAHE7B	11	121.57			
C	Parupeneus multifasciatus	9	KAHE7B	1	96.08			
C	Bodianus bilunulatus	9	KAHE7B	1	0.52			
C	Thalassoma duperrey	9	KAHE7B	1	97.05			
C	Thalassoma duperrey	9	KAHE7B	1	54.95			
C	Sufflamen bursa	9	KAHE7B	1	144.65			
C	Sufflamen bursa	9	KAHE7B	1	85.87	97	1229.19	77.5
H	Acanthurus nigrofuscus	9	KAHE7B	1	54.25	1	54.25	3.4
O	Melichthys vidua	9	KAHE7B	1	296.24			
O	Canthigaster jactator	9	KAHE7B	1	2.21			
O	Canthigaster jactator	9	KAHE7B	1	3.56	3	302.01	19.0
P	Chromis vanderbilti	9	KAHE7B	5	1.58	5	1.58	0.1
	TOTAL	9	KAHE7B	106	1587.02	106	1587.02	100.000001

14-Jun-13						GROUP		GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	10	KAHE7C	1	3.12			
C	Parupeneus multifasciatus	10	KAHE7C	1	96.08			
C	Plectroglyphidodon johnstonianus	10	KAHE7C	1	1.72			
C	Paracirrhites arcatus	10	KAHE7C	2	32.70			
C	Thalassoma duperrey	10	KAHE7C	3	9.45			
C	Sufflamen bursa	10	KAHE7C	5	723.25			
C	Sufflamen bursa	10	KAHE7C	2	171.75	15	1038.07	68.5
O	Melichthys vidua	10	KAHE7C	1	296.24			
O	Cantherhines sandwichiensis	10	KAHE7C	2	164.10			
O	Canthigaster coronata	10	KAHE7C	1	7.59			
O	Canthigaster jactator	10	KAHE7C	2	7.12	6	475.06	31.3
P	Chromis vanderbilti	10	KAHE7C	7	2.21	7	2.21	0.1
	TOTAL	10	KAHE7C	28	1515.33	28	1515.33	100
C	Parupeneus pleurostigma	11	KAHE7D	1	4.11			
C	Parupeneus multifasciatus	11	KAHE7D	1	3.12			
C	Coris gaimard	11	KAHE7D	1	0.23			
C	Sufflamen bursa	11	KAHE7D	1	85.87			
C	Sufflamen fraenatus	11	KAHE7D	1	224.79	5	318.12	97.8
O	Canthigaster jactator	11	KAHE7D	2	7.12	2	7.12	2.2
	TOTAL	11	KAHE7D	7	325.25	7	325.25	100.000001
C	Fistularia commersoni	12	KAHE7E	1	45.57			
C	Parupeneus pleurostigma	12	KAHE7E	107	439.37			
C	Parupeneus pleurostigma	12	KAHE7E	1	58.01			
C	Parupeneus multifasciatus	12	KAHE7E	173	539.59			
C	Parupeneus multifasciatus	12	KAHE7E	1	235.75			
C	Paracirrhites arcatus	12	KAHE7E	3	10.34			
C	Labroides phthiophagus	12	KAHE7E	1	0.19			
C	Cheilinus bimaculatus	12	KAHE7E	1	9.63			
C	Pseudocheilinus octotaenia	12	KAHE7E	1	4.00			
C	Thalassoma duperrey	12	KAHE7E	1	97.05			
C	Thalassoma duperrey	12	KAHE7E	2	54.78			
C	Thalassoma duperrey	12	KAHE7E	1	54.95			
C	Pseudojuloides cerasinus	12	KAHE7E	3	3.85			
C	Sufflamen bursa	12	KAHE7E	4	343.49	300	1896.57	27.0
CF	Chaetodon multicinctus	12	KAHE7E	1	2.80	1	2.80	0.0
H	Acanthurus nigrofuscus	12	KAHE7E	3	22.57			
H	Acanthurus nigrofuscus	12	KAHE7E	17	17.74			
H	Acanthurus olivaceus	12	KAHE7E	1	261.67			
H	Acanthurus olivaceus	12	KAHE7E	1	393.21			
H	Zebrasoma flavescens	12	KAHE7E	1	1.69			
H	Naso lituratus	12	KAHE7E	2	145.17			
H	Naso lituratus	12	KAHE7E	1	205.99			
H	Naso unicornis	12	KAHE7E	1	38.92	27	1086.96	15.5
O	Melichthys niger	12	KAHE7E	18	2940.04			
O	Melichthys vidua	12	KAHE7E	2	398.06			
O	Canthigaster jactator	12	KAHE7E	2	7.12	22	3345.22	47.6
P	Chaetodon miliaris	12	KAHE7E	1	0.88			
P	Chromis vanderbilti	12	KAHE7E	52	16.43			
P	Naso brevirostris	12	KAHE7E	4	293.43			
P	Naso brevirostris	12	KAHE7E	2	381.49	59	692.23	9.9
	TOTAL	12	KAHE7E	409	7023.79	409	7023.79	99.9999998
C	Lutjanus kasmira	13	KAHE10	149	16383.85			
C	Mulloides flavolineatus	13	KAHE10	31	8349.57			
C	Parupeneus pleurostigma	13	KAHE10	1	30.44			
C	Parupeneus multifasciatus	13	KAHE10	3	81.35			
C	Parupeneus multifasciatus	13	KAHE10	4	384.33			
C	Parupeneus multifasciatus	13	KAHE10	133	414.83			
C	Parupeneus multifasciatus	13	KAHE10	4	217.60			
C	Paracirrhites arcatus	13	KAHE10	1	8.12			
C	Labroides phthiophagus	13	KAHE10	2	1.26			
C	Thalassoma duperrey	13	KAHE10	7	191.73			
C	Thalassoma duperrey	13	KAHE10	10	549.49			
C	Thalassoma duperrey	13	KAHE10	4	44.65			
C	Thalassoma duperrey	13	KAHE10	4	388.21			
C	Stethojulis balteata	13	KAHE10	1	72.39			
C	Stethojulis balteata	13	KAHE10	2	71.53			
C	Macropharyngodon geoffroy	13	KAHE10	1	18.63			
C	Halichoeres ornatissimus	13	KAHE10	1	16.45			
C	Sufflamen bursa	13	KAHE10	2	171.75	360	27396.16	94.4
CF	Chaetodon multicinctus	13	KAHE10	1	13.03	1	13.03	0.04
H	Acanthurus triostegus	13	KAHE10	3	51.50			
H	Acanthurus triostegus	13	KAHE10	3	138.95			
H	Acanthurus nigrofuscus	13	KAHE10	3	162.76			
H	Acanthurus nigrofuscus	13	KAHE10	1	14.21			
H	Acanthurus nigrofuscus	13	KAHE10	6	143.38			
H	Acanthurus blochii	13	KAHE10	1	29.03	17	539.83	1.9
O	Stegastes fasciatus	13	KAHE10	10	259.80			
O	Cantherhines sandwichiensis	13	KAHE10	2	164.10			
O	Canthigaster coronata	13	KAHE10	1	7.59			
O	Canthigaster jactator	13	KAHE10	1	3.56	14	435.06	1.5
P	Dascyllus albisella	13	KAHE10	12	37.06			
P	Abudefduf abdominalis	13	KAHE10	10	313.33			
P	Chromis vanderbilti	13	KAHE10	31	9.79			
P	Chromis ovalis	13	KAHE10	23	265.87	76	626.05	2.2
	TOTAL	13	KAHE10	468	29010.12	468	29010.12	100

14-Jun-13						GROUP		GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus pleurostigma	14	NANA1	3	12.32			
C	Parupeneus multifasciatus	14	NANA1	18	56.14			
C	Plectroglyphidodon imparipennis	14	NANA1	6	5.17			
C	Thalassoma lutescens	14	NANA1	1	54.95			
C	Thalassoma duperrey	14	NANA1	8	25.20			
C	Thalassoma duperrey	14	NANA1	5	55.82			
C	Thalassoma duperrey	14	NANA1	4	109.56			
C	Rhinecanthus rectangulus	14	NANA1	3	257.62			
C	Rhinecanthus rectangulus	14	NANA1	1	45.36	49	622.15	88.0
H	Acanthurus triostegus	14	NANA1	1	46.32			
H	Acanthurus triostegus	14	NANA1	2	34.33	3	80.65	11.4
O	Canthigaster jactator	14	NANA1	1	3.56	1	3.56	0.5
P	Chromis vanderbilti	14	NANA1	3	0.95	3	0.95	0.1
	TOTAL	14	NANA1	56	707.31	56	707.31	100
C	Parupeneus multifasciatus	15	NANA2	7	21.83			
C	Parupeneus multifasciatus	15	NANA2	1	96.08			
C	Labroides phthirophagus	15	NANA2	1	0.63			
C	Thalassoma duperrey	15	NANA2	3	82.17			
C	Thalassoma duperrey	15	NANA2	5	274.74			
C	Thalassoma duperrey	15	NANA2	6	582.31			
C	Gomphosus varius	15	NANA2	2	22.08			
C	Gomphosus varius	15	NANA2	1	39.39			
C	Halichoeres ornatissimus	15	NANA2	1	25.14			
C	Zanclus cornutus	15	NANA2	4	416.63	31	1561.02	9.6
CF	Chaetodon multicinctus	15	NANA2	2	26.06			
CF	Chaetodon multicinctus	15	NANA2	2	13.31	4	39.37	0.2
H	Acanthurus triostegus	15	NANA2	3	300.04			
H	Acanthurus triostegus	15	NANA2	2	92.63			
H	Acanthurus leucopareius	15	NANA2	10	1327.00			
H	Acanthurus leucopareius	15	NANA2	8	1861.40			
H	Acanthurus leucopareius	15	NANA2	5	333.70			
H	Acanthurus nigrofuscus	15	NANA2	14	198.98			
H	Acanthurus nigrofuscus	15	NANA2	15	358.46			
H	Acanthurus olivaceus	15	NANA2	15	8447.55			
H	Acanthurus blochii	15	NANA2	3	696.72			
H	Ctenochaetus strigosus	15	NANA2	2	53.69	77	13670.18	84.4
O	Stegastes fasciolatus	15	NANA2	7	103.34			
O	Melichthys niger	15	NANA2	5	816.68	12	920.02	5.7
P	Chromis vanderbilti	15	NANA2	43	13.58	43	13.58	0.1
	TOTAL	15	NANA2	167	16204.17	167	16204.17	99.9999998
C	Aulostomus chinensis	16	PIPE	1	52.00			
C	Decapterus macarellus	16	PIPE	35	4559.77			
C	Aprion virescens	16	PIPE	1	2267.96			
C	Lutjanus kasmira	16	PIPE	15	1649.38			
C	Mulloides flavolineatus	16	PIPE	1	130.28			
C	Parupeneus multifasciatus	16	PIPE	3	288.24			
C	Parupeneus cyclostomus	16	PIPE	1	88.57			
C	Forcipiger flavissimus	16	PIPE	1	5.13			
C	Plectroglyphidodon johnstonianus	16	PIPE	3	5.17			
C	Pseudocheilinus octotaenia	16	PIPE	3	107.29			
C	Thalassoma duperrey	16	PIPE	14	383.47			
C	Thalassoma duperrey	16	PIPE	21	1153.93			
C	Thalassoma duperrey	16	PIPE	7	78.14			
C	Thalassoma duperrey	16	PIPE	51	4949.65			
C	Gomphosus varius	16	PIPE	3	118.18			
C	Gomphosus varius	16	PIPE	4	90.40			
C	Coris venusta	16	PIPE	2	96.76			
C	Coris venusta	16	PIPE	3	70.91			
C	Halichoeres ornatissimus	16	PIPE	3	49.35			
C	Zanclus cornutus	16	PIPE	1	104.16			
C	Sufflamen bursa	16	PIPE	1	85.87	174	16334.62	27.8
CF	Chaetodon multicinctus	16	PIPE	4	52.12	4	52.12	0.1
H	Calotomus carolinus	16	PIPE	1	707.00			
H	Scarus sordidus	16	PIPE	27	2052.36			
H	Scarus sordidus	16	PIPE	14	1960.08			
H	Scarus sordidus	16	PIPE	25	899.87			
H	Scarus sordidus	16	PIPE	5	7137.85			
H	Scarus sordidus	16	PIPE	11	4037.24			
H	Scarus psittacus	16	PIPE	2	1572.31			
H	Scarus psittacus	16	PIPE	3	1664.07			
H	Scarus psittacus	16	PIPE	3	4312.39			
H	Acanthurus nigrofuscus	16	PIPE	17	922.29			
H	Acanthurus nigrofuscus	16	PIPE	27	645.22			
H	Acanthurus blochii	16	PIPE	3	293.93			
H	Ctenochaetus strigosus	16	PIPE	6	161.06			
H	Naso lituratus	16	PIPE	3	1346.83	147	27712.49	47.2
O	Stegastes fasciolatus	16	PIPE	25	369.08			
O	Melichthys niger	16	PIPE	1	99.66			
O	Melichthys vidua	16	PIPE	1	199.03			
O	Cantherhines sandwichiensis	16	PIPE	1	82.05	28	749.82	1.3
P	Chaetodon miliaris	16	PIPE	2	42.34			
P	Abudefduf abdominalis	16	PIPE	92	2882.60			
P	Abudefduf vaigensis	16	PIPE	196	9182.97			
P	Chromis vanderbilti	16	PIPE	19	6.00			
P	Chromis ovalis	16	PIPE	28	166.09			
P	Chromis ovalis	16	PIPE	123	1421.84			
P	Naso brevirostris	16	PIPE	3	220.07	463	13921.91	23.7
	TOTAL	16	PIPE	816	58770.96	816	58770.96	100

20 SEPTEMBER 2013 FIELD DATA

20-Sep-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	1	EAST1	4	217.60			
C	Parupeneus multifasciatus	1	EAST1	1	340.44			
C	Parupeneus multifasciatus	1	EAST1	3	81.35			
C	Parupeneus cyclostomus	1	EAST1	1	24.80			
C	Parupeneus cyclostomus	1	EAST1	1	88.57			
C	Parupeneus cyclostomus	1	EAST1	1	49.96			
C	Plectroglyphidodon johnstonianus	1	EAST1	2	3.44			
C	Plectroglyphidodon imparipennis	1	EAST1	2	1.72			
C	Paracirrhites arcatus	1	EAST1	4	32.47			
C	Paracirrhites forsteri	1	EAST1	1	39.65			
C	Bodianus bilunulatus	1	EAST1	1	1141.37			
C	Thalassoma duperrey	1	EAST1	11	301.30			
C	Thalassoma duperrey	1	EAST1	18	56.71			
C	Thalassoma duperrey	1	EAST1	8	439.59			
C	Thalassoma duperrey	1	EAST1	3	291.16			
C	Thalassoma duperrey	1	EAST1	12	133.96			
C	Halichoeres ornatissimus	1	EAST1	1	16.45			
C	Halichoeres ornatissimus	1	EAST1	1	25.14			
C	Rhinecanthus rectangulus	1	EAST1	1	45.36	76	3331.04	18.5
H	Calotomus carolinus	1	EAST1	1	3318.76			
H	Acanthurus triostegus	1	EAST1	3	51.50			
H	Acanthurus triostegus	1	EAST1	9	416.84			
H	Acanthurus triostegus	1	EAST1	15	1500.21			
H	Acanthurus triostegus	1	EAST1	1	1.57			
H	Acanthurus nigrofuscus	1	EAST1	9	67.72			
H	Acanthurus nigrofuscus	1	EAST1	7	379.77			
H	Acanthurus nigrofuscus	1	EAST1	19	454.04			
H	Acanthurus olivaceus	1	EAST1	8	2093.34			
H	Acanthurus olivaceus	1	EAST1	9	5068.53			
H	Acanthurus olivaceus	1	EAST1	3	142.43			
H	Acanthurus blochii	1	EAST1	1	783.81			
H	Naso lituratus	1	EAST1	1	72.58	86	14351.11	79.6
O	Stegastes fasciolatus	1	EAST1	1	14.76			
O	Melichthys vidua	1	EAST1	1	296.24			
O	Canthigaster jactator	1	EAST1	5	17.81	7	328.81	1.8
P	Heniochus diphreutes	1	EAST1	1	14.25			
P	Chromis vanderbilti	1	EAST1	36	11.37	37	25.62	0.1
	TOTAL	1	EAST1	206	18036.58	206	18036.58	100
C	Parupeneus multifasciatus	2	EAST2	4	44.21			
C	Parupeneus multifasciatus	2	EAST2	4	108.47			
C	Plectroglyphidodon johnstonianus	2	EAST2	1	1.72			
C	Plectroglyphidodon imparipennis	2	EAST2	1	0.86			
C	Bodianus bilunulatus	2	EAST2	1	324.75			
C	Thalassoma duperrey	2	EAST2	11	122.80			
C	Thalassoma duperrey	2	EAST2	4	388.21			
C	Thalassoma duperrey	2	EAST2	12	328.69			
C	Thalassoma duperrey	2	EAST2	6	329.69			
C	Coris gaimard	2	EAST2	1	8.57			
C	Stethojulis balteata	2	EAST2	3	107.29			
C	Sufflamen fraenatus	2	EAST2	1	85.87	49	1851.12	16.8
CF	Chaetodon quadrimaculatus	2	EAST2	1	25.30			
CF	Chaetodon multicinctus	2	EAST2	2	1.65	3	26.96	0.2
H	Scarus sordidus	2	EAST2	1	13.73			
H	Scarus psittacus	2	EAST2	2	7.64			
H	Acanthurus nigrofuscus	2	EAST2	1	54.25			
H	Acanthurus nigrofuscus	2	EAST2	18	255.83			
H	Acanthurus nigrofuscus	2	EAST2	10	75.25			
H	Acanthurus nigrofuscus	2	EAST2	16	382.35			
H	Acanthurus olivaceus	2	EAST2	15	5898.15			
H	Acanthurus olivaceus	2	EAST2	1	563.17			
H	Acanthurus olivaceus	2	EAST2	7	1144.61			
H	Acanthurus blochii	2	EAST2	1	453.59			
H	Zebrasoma flavescens	2	EAST2	1	53.28	73	8901.86	81.0
O	Melichthys vidua	2	EAST2	1	199.03			
O	Canthigaster jactator	2	EAST2	3	10.69	4	209.72	1.9
	TOTAL	2	EAST2	129	10989.65	129	10989.65	100

20-Sep-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Parupeneus multifasciatus	3	EAST3	2	54.23			
C	Parupeneus multifasciatus	3	EAST3	3	163.20			
C	Plectroglyphidodon johnstonianus	3	EAST3	5	8.61			
C	Paracirrhites arcatus	3	EAST3	3	24.35			
C	Paracirrhites arcatus	3	EAST3	3	49.05			
C	Pseudocheilinus octotaenia	3	EAST3	1	14.41			
C	Thalassoma duperrey	3	EAST3	5	136.95			
C	Thalassoma duperrey	3	EAST3	5	55.82			
C	Thalassoma duperrey	3	EAST3	7	384.64			
C	Coris gaimard	3	EAST3	3	66.21			
C	Sufflamen bursa	3	EAST3	5	429.37			
C	Sufflamen fraenatus	3	EAST3	1	144.65	43	1531.49	48.6
CF	Chaetodon unimaculatus	3	EAST3	1	2.83			
CF	Chaetodon unimaculatus	3	EAST3	5	126.52			
CF	Chaetodon ornatissimus	3	EAST3	2	138.00			
CF	Chaetodon multicinctus	3	EAST3	3	2.48			
CF	Chaetodon multicinctus	3	EAST3	2	26.06	13	295.89	9.4
H	Acanthurus nigrofuscus	3	EAST3	11	82.77			
H	Acanthurus nigrofuscus	3	EAST3	11	262.87			
H	Acanthurus nigrofuscus	3	EAST3	17	241.62			
H	Acanthurus blochii	3	EAST3	1	56.70			
H	Ctenochaetus strigosus	3	EAST3	8	24.70			
H	Ctenochaetus strigosus	3	EAST3	4	107.38			
H	Ctenochaetus strigosus	3	EAST3	5	75.99			
H	Ctenochaetus strigosus	3	EAST3	1	0.87			
H	Zebrasoma flavescens	3	EAST3	2	3.38			
H	Zebrasoma flavescens	3	EAST3	1	26.03	61	882.30	28.0
O	Stegastes fasciatus	3	EAST3	3	44.29			
O	Melichthys niger	3	EAST3	1	163.34			
O	Melichthys vidua	3	EAST3	1	199.03			
O	Canthigaster jactator	3	EAST3	10	35.62	15	442.28	14.0
	TOTAL	3	EAST3	132	3151.96	132	3151.96	100
C	Parupeneus pleurostigma	4	EAST4	2	8.21			
C	Parupeneus pleurostigma	4	EAST4	3	91.31			
C	Parupeneus pleurostigma	4	EAST4	4	53.02			
C	Parupeneus multifasciatus	4	EAST4	5	135.58			
C	Parupeneus multifasciatus	4	EAST4	1	96.08			
C	Plectroglyphidodon johnstonianus	4	EAST4	1	1.72			
C	Paracirrhites arcatus	4	EAST4	1	16.35			
C	Paracirrhites arcatus	4	EAST4	2	16.24			
C	Cirrhitops fasciatus	4	EAST4	1	3.75			
C	Bodianus bilunulatus	4	EAST4	1	1141.37			
C	Bodianus bilunulatus	4	EAST4	1	324.75			
C	Labroides phthirophagus	4	EAST4	1	0.63			
C	Pseudocheilinus octotaenia	4	EAST4	1	35.76			
C	Thalassoma duperrey	4	EAST4	7	384.64			
C	Thalassoma duperrey	4	EAST4	15	410.86			
C	Thalassoma duperrey	4	EAST4	17	189.78			
C	Coris venusta	4	EAST4	2	47.27			
C	Coris gaimard	4	EAST4	1	8.57			
C	Pseudojuloides cerasinus	4	EAST4	2	22.33			
C	Stethojulis balteata	4	EAST4	3	43.23			
C	Halichoeres ornatissimus	4	EAST4	3	75.41			
C	Halichoeres ornatissimus	4	EAST4	3	28.57			
C	Rhinecanthus rectangulus	4	EAST4	4	181.45			
C	Sufflamen fraenatus	4	EAST4	1	144.65			
C	Sufflamen fraenatus	4	EAST4	1	85.87	83	3547.40	72.4
H	Acanthurus nigrofuscus	4	EAST4	20	284.25			
H	Acanthurus nigrofuscus	4	EAST4	9	215.07			
H	Acanthurus nigrofuscus	4	EAST4	15	112.87			
H	Ctenochaetus strigosus	4	EAST4	1	3.09	45	615.28	12.6
O	Melichthys niger	4	EAST4	1	163.34			
O	Melichthys vidua	4	EAST4	2	398.06			
O	Melichthys vidua	4	EAST4	1	124.35			
O	Canthigaster jactator	4	EAST4	5	17.81	9	703.55	14.4
P	Chromis vanderbilti	4	EAST4	103	32.54	103	32.54	0.7
	TOTAL	4	EAST4	240	4898.77	240	4898.77	100

20-Sep-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Monotaxis grandoculis	5	KO1	3	104.37				
C	Monotaxis grandoculis	5	KO1	3	214.58				
C	Monotaxis grandoculis	5	KO1	1	212.06				
C	Parupeneus multifasciatus	5	KO1	4	108.47				
C	Parupeneus multifasciatus	5	KO1	2	108.80				
C	Parupeneus cyclostomus	5	KO1	1	24.80				
C	Plectroglyphidodon johnstonianus	5	KO1	1	1.72				
C	Labroides phthiophagus	5	KO1	1	0.63				
C	Thalassoma duperrey	5	KO1	13	714.34				
C	Thalassoma duperrey	5	KO1	15	410.86				
C	Thalassoma duperrey	5	KO1	9	873.47				
C	Gomphosus varius	5	KO1	2	22.08				
C	Gomphosus varius	5	KO1	1	62.03				
C	Stethojulis balteata	5	KO1	10	357.63				
C	Stethojulis balteata	5	KO1	4	289.56				
C	Zanclus cornutus	5	KO1	1	104.16				
C	Sufflamen bursa	5	KO1	2	171.75				
C	Sufflamen bursa	5	KO1	1	144.65	74	3925.94	21.8	
CF	Chaetodon unimaculatus	5	KO1	2	50.61				
CF	Chaetodon ornatissimus	5	KO1	5	345.01				
CF	Chaetodon quadrimaculatus	5	KO1	2	50.61				
CF	Chaetodon multicinctus	5	KO1	6	78.18	15	524.41	2.9	
H	Scarus sordidus	5	KO1	2	469.30				
H	Scarus sordidus	5	KO1	1	35.99				
H	Scarus sordidus	5	KO1	2	152.03				
H	Acanthurus triostegus	5	KO1	42	1945.23				
H	Acanthurus triostegus	5	KO1	2	18.30				
H	Acanthurus nigrofuscus	5	KO1	24	573.53				
H	Acanthurus nigrofuscus	5	KO1	9	488.27				
H	Acanthurus olivaceus	5	KO1	8	1308.13				
H	Acanthurus olivaceus	5	KO1	14	1312.76				
H	Acanthurus olivaceus	5	KO1	2	523.33				
H	Acanthurus blochii	5	KO1	5	145.15				
H	Acanthurus blochii	5	KO1	8	453.59				
H	Ctenochaetus strigosus	5	KO1	40	1073.76				
H	Ctenochaetus strigosus	5	KO1	5	660.67				
H	Ctenochaetus strigosus	5	KO1	44	2898.06				
H	Zebrasoma flavescens	5	KO1	1	1.69				
H	Zebrasoma flavescens	5	KO1	6	156.18				
H	Zebrasoma flavescens	5	KO1	4	37.94	219	12253.92	67.9	
O	Stegastes fasciolatus	5	KO1	2	29.53				
O	Melichthys niger	5	KO1	6	980.01				
O	Melichthys vidua	5	KO1	1	199.03				
O	Canthigaster jactator	5	KO1	3	6.62	12	1215.19	6.7	
P	Abudefduf abdominalis	5	KO1	4	125.33	4	125.33	0.7	
	TOTAL	5	KO1	324	18044.78	324	18044.78	100	
C	Cephalopholis argus	6	KO2	1	1676.50				
C	Cephalopholis argus	6	KO2	1	229.32				
C	Parupeneus pleurostigma	6	KO2	1	30.44				
C	Parupeneus multifasciatus	6	KO2	1	96.08				
C	Parupeneus multifasciatus	6	KO2	1	235.75				
C	Parupeneus multifasciatus	6	KO2	2	54.23				
C	Parupeneus multifasciatus	6	KO2	2	108.80				
C	Parupeneus bifasciatus	6	KO2	1	11.60				
C	Parupeneus cyclostomus	6	KO2	1	10.05				
C	Parupeneus cyclostomus	6	KO2	1	49.96				
C	Parupeneus cyclostomus	6	KO2	1	24.80				
C	Forcipiger flavissimus	6	KO2	1	9.15				
C	Chaetodon lunula	6	KO2	2	71.99				
C	Plectroglyphidodon johnstonianus	6	KO2	3	5.17				
C	Bodianus bilunulatus	6	KO2	1	1141.37				
C	Bodianus bilunulatus	6	KO2	1	214.67				
C	Labroides phthiophagus	6	KO2	3	1.88				
C	Thalassoma duperrey	6	KO2	8	439.59				
C	Thalassoma duperrey	6	KO2	7	679.36				
C	Thalassoma duperrey	6	KO2	9	246.51				
C	Stethojulis balteata	6	KO2	2	144.78				
C	Stethojulis balteata	6	KO2	2	71.53				
C	Sufflamen bursa	6	KO2	2	171.75				
C	Ostracion meleagris	6	KO2	1	6.76	55	5732.04	22.7	
CF	Chaetodon unimaculatus	6	KO2	6	151.82				
CF	Chaetodon quadrimaculatus	6	KO2	6	151.82				
CF	Chaetodon multicinctus	6	KO2	6	78.18	18	381.83	1.5	
H	Scarus psittacus	6	KO2	1	1437.46				
H	Scarus psittacus	6	KO2	1	554.69				
H	Scarus psittacus	6	KO2	12	454.47				
H	Scarus psittacus	6	KO2	11	160.76				
H	Acanthurus triostegus	6	KO2	14	648.41				
H	Acanthurus leucopareius	6	KO2	1	132.70				
H	Acanthurus leucopareius	6	KO2	1	66.74				
H	Acanthurus nigrofuscus	6	KO2	6	325.51				
H	Acanthurus nigrofuscus	6	KO2	13	310.66				
H	Acanthurus olivaceus	6	KO2	12	6758.04				
H	Acanthurus olivaceus	6	KO2	2	187.54				
H	Acanthurus olivaceus	6	KO2	6	981.10				
H	Acanthurus blochii	6	KO2	1	29.03				
H	Acanthurus blochii	6	KO2	2	311.17				
H	Acanthurus blochii	6	KO2	2	113.40				
H	Ctenochaetus strigosus	6	KO2	7	53.03				
H	Ctenochaetus strigosus	6	KO2	6	792.80				
H	Ctenochaetus strigosus	6	KO2	4	263.46				
H	Ctenochaetus strigosus	6	KO2	8	214.75				
H	Ctenochaetus strigosus	6	KO2	9	7.84				
H	Zebrasoma flavescens	6	KO2	11	104.33				
H	Zebrasoma flavescens	6	KO2	11	586.08				
H	Zebrasoma flavescens	6	KO2	9	234.27				
H	Naso lituratus	6	KO2	2	1244.71				
H	Naso lituratus	6	KO2	1	205.99	153	16178.94	64.2	
O	Melichthys niger	6	KO2	13	2123.36				
O	Melichthys vidua	6	KO2	1	199.03				
O	Cantherhines sandwichiensis	6	KO2	1	82.05				
O	Canthigaster jactator	6	KO2	2	7.12	17	2411.57	9.6	
P	Abudefduf abdominalis	6	KO2	16	501.32	16	501.32	2.0	
	TOTAL	6	KO2	259	25205.71	259	25205.71	100	

20-Sep-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Parupeneus multifasciatus	7	KAHE1D	2	54.23				
C	Parupeneus bifasciatus	7	KAHE1D	1	107.31				
C	Parupeneus bifasciatus	7	KAHE1D	1	29.20				
C	Plectroglyphidodon johnstonianus	7	KAHE1D	5	8.61				
C	Plectroglyphidodon imparipennis	7	KAHE1D	1	0.86				
C	Paracirrhites arcatus	7	KAHE1D	1	8.12				
C	Labroides phthirophagus	7	KAHE1D	1	0.63				
C	Thalassoma duperrey	7	KAHE1D	17	1649.88				
C	Thalassoma duperrey	7	KAHE1D	7	191.73				
C	Thalassoma duperrey	7	KAHE1D	47	2582.60				
C	Gomphosus varius	7	KAHE1D	5	55.21				
C	Stethojulis balteata	7	KAHE1D	5	178.82				
C	Stethojulis balteata	7	KAHE1D	6	86.45				
C	Stethojulis balteata	7	KAHE1D	3	217.17				
C	Halichoeres ornatissimus	7	KAHE1D	1	25.14				
C	Halichoeres ornatissimus	7	KAHE1D	1	16.45				
C	Sufflamen bursa	7	KAHE1D	2	171.75	106	5384.15	28.3	
CF	Chaetodon quadrimaculatus	7	KAHE1D	3	75.91				
CF	Chaetodon multicinctus	7	KAHE1D	4	52.12	7	128.03	0.7	
H	Scarus sordidus	7	KAHE1D	2	152.03				
H	Scarus psittacus	7	KAHE1D	1	37.87				
H	Acanthurus triostegus	7	KAHE1D	3	138.95				
H	Acanthurus leucopareus	7	KAHE1D	1	15.69				
H	Acanthurus leucopareus	7	KAHE1D	1	7.89				
H	Acanthurus nigrofusus	7	KAHE1D	5	71.06				
H	Acanthurus nigrofusus	7	KAHE1D	4	217.01				
H	Acanthurus nigrofusus	7	KAHE1D	22	525.73				
H	Acanthurus olivaceus	7	KAHE1D	5	237.38				
H	Acanthurus olivaceus	7	KAHE1D	11	217.18				
H	Acanthurus blochii	7	KAHE1D	1	155.58				
H	Ctenochaetus strigosus	7	KAHE1D	10	268.44				
H	Ctenochaetus strigosus	7	KAHE1D	7	53.03				
H	Ctenochaetus strigosus	7	KAHE1D	5	329.32				
H	Zebbrasoma flavescens	7	KAHE1D	2	106.56	80	2533.74	13.3	
O	Stegastes fasciolatus	7	KAHE1D	8	118.11				
O	Melichthys niger	7	KAHE1D	63	10290.14				
O	Melichthys vidua	7	KAHE1D	1	199.03				
O	Canthigaster jactator	7	KAHE1D	3	6.62	75	10613.89	55.9	
P	Abudefduf vaigensis	7	KAHE1D	9	281.99				
P	Chromis ovalis	7	KAHE1D	5	57.80	14	339.79	1.8	
	TOTAL	7	KAHE1D	282	18999.60	282	18999.60	100	
C	Parupeneus pleurostigma	8	KAHE5B	1	30.44				
C	Parupeneus multifasciatus	8	KAHE5B	1	11.05				
C	Parupeneus multifasciatus	8	KAHE5B	6	162.70				
C	Parupeneus multifasciatus	8	KAHE5B	3	288.24				
C	Parupeneus cyclostomus	8	KAHE5B	1	49.96				
C	Parupeneus cyclostomus	8	KAHE5B	1	10.05				
C	Plectroglyphidodon johnstonianus	8	KAHE5B	1	1.72				
C	Paracirrhites arcatus	8	KAHE5B	3	24.35				
C	Cirrhites fasciatus	8	KAHE5B	1	8.23				
C	Novaculichthys taeniourus	8	KAHE5B	1	64.38				
C	Thalassoma duperrey	8	KAHE5B	5	274.74				
C	Thalassoma duperrey	8	KAHE5B	4	388.21				
C	Thalassoma duperrey	8	KAHE5B	2	22.33				
C	Thalassoma duperrey	8	KAHE5B	2	54.78				
C	Gomphosus varius	8	KAHE5B	1	62.03				
C	Gomphosus varius	8	KAHE5B	1	22.60				
C	Coris venusta	8	KAHE5B	1	23.64				
C	Stethojulis balteata	8	KAHE5B	1	72.39				
C	Halichoeres ornatissimus	8	KAHE5B	1	25.14				
C	Zanclus cornutus	8	KAHE5B	3	72.13				
C	Zanclus cornutus	8	KAHE5B	3	312.47				
C	Sufflamen bursa	8	KAHE5B	1	85.87	44	2067.45	15.8	
CF	Chaetodon unimaculatus	8	KAHE5B	1	25.30				
CF	Chaetodon unimaculatus	8	KAHE5B	1	0.79				
CF	Chaetodon multicinctus	8	KAHE5B	4	52.12	6	78.21	0.6	
H	Calotomus carolinus	8	KAHE5B	1	7.39				
H	Acanthurus triostegus	8	KAHE5B	1	9.15				
H	Acanthurus nigrofusus	8	KAHE5B	12	170.55				
H	Acanthurus nigrofusus	8	KAHE5B	11	596.78				
H	Acanthurus nigrofusus	8	KAHE5B	10	75.25				
H	Acanthurus olivaceus	8	KAHE5B	6	1570.00				
H	Acanthurus olivaceus	8	KAHE5B	14	7884.38				
H	Acanthurus olivaceus	8	KAHE5B	1	93.77				
H	Naso lituratus	8	KAHE5B	2	3.47	58	10410.75	79.8	
O	Stegastes fasciolatus	8	KAHE5B	5	73.82				
O	Melichthys vidua	8	KAHE5B	1	296.24				
O	Cantherhines sandwichiensis	8	KAHE5B	1	46.04				
O	Canthigaster jactator	8	KAHE5B	4	14.25	11	430.34	3.3	
P	Chromis vanderbilti	8	KAHE5B	64	20.22				
P	Naso brevirostris	8	KAHE5B	1	38.92	65	59.14	0.5	
	TOTAL	8	KAHE5B	184	13045.89	184	13045.89	100	
C	Hyporhamphus pacificus	9	KAHE7B	180	317070.89				
C	Aulostomus chinensis	9	KAHE7B	1	12.84				
C	Mulloides flavolineatus	9	KAHE7B	13	4747.72				
C	Parupeneus multifasciatus	9	KAHE7B	2	192.16				
C	Parupeneus multifasciatus	9	KAHE7B	2	108.80				
C	Parupeneus multifasciatus	9	KAHE7B	1	155.42				
C	Parupeneus multifasciatus	9	KAHE7B	4	44.21				
C	Parupeneus multifasciatus	9	KAHE7B	3	81.35				
C	Pseudocheilinus octotaenia	9	KAHE7B	1	1.61				
C	Thalassoma duperrey	9	KAHE7B	2	194.10				
C	Thalassoma duperrey	9	KAHE7B	2	109.90				
C	Coris venusta	9	KAHE7B	1	9.39				
C	Sufflamen bursa	9	KAHE7B	2	171.75				
C	Sufflamen bursa	9	KAHE7B	2	289.30	216	323189.43	99.8	
CF	Chaetodon multicinctus	9	KAHE7B	1	6.66				
CF	Chaetodon multicinctus	9	KAHE7B	3	2.48	4	9.14	0.003	
H	Acanthurus triostegus	9	KAHE7B	1	17.17				
H	Acanthurus nigrofusus	9	KAHE7B	2	108.50				
H	Acanthurus nigrofusus	9	KAHE7B	4	4.17				
H	Naso unicornis	9	KAHE7B	3	16.31				
H	Naso unicornis	9	KAHE7B	1	190.74	11	336.90	0.1	
O	Melichthys vidua	9	KAHE7B	1	199.03				
O	Cantherhines sandwichiensis	9	KAHE7B	1	82.05				
O	Canthigaster jactator	9	KAHE7B	5	17.81	7	298.89	0.1	
P	Chaetodon miliaris	9	KAHE7B	1	0.88				
P	Chromis vanderbilti	9	KAHE7B	27	8.53	28	9.41	0.003	
	TOTAL	9	KAHE7B	266	323843.77	266	323843.77	100	

20-Sep-13						GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS PERCENT
C	Parupeneus multifasciatus	10	KAHE7C	1	54.40		
C	Parupeneus multifasciatus	10	KAHE7C	2	680.89		
C	Parupeneus multifasciatus	10	KAHE7C	4	108.47		
C	Parupeneus cyclostomus	10	KAHE7C	1	49.96		
C	Paracirrhites arcatus	10	KAHE7C	1	3.45		
C	Paracirrhites arcatus	10	KAHE7C	2	16.24		
C	Gomphosus varius	10	KAHE7C	1	91.05		
C	Zanclus cornutus	10	KAHE7C	1	104.16		
C	Sufflamen bursa	10	KAHE7C	4	343.49		
C	Sufflamen bursa	10	KAHE7C	4	578.60	21	2030.70 86.8
CF	Chaetodon multicinctus	10	KAHE7C	1	0.83		
CF	Pervagor spilosoma	10	KAHE7C	1	16.22	2	17.04 0.7
H	Naso lituratus	10	KAHE7C	1	1.74		
H	Naso unicornis	10	KAHE7C	8	19.21	9	20.95 0.9
O	Melichthys vidua	10	KAHE7C	1	199.03		
O	Cantherhines sandwichiensis	10	KAHE7C	1	46.04		
O	Canthigaster coronata	10	KAHE7C	1	13.65		
O	Canthigaster jactator	10	KAHE7C	2	7.12	5	265.84 11.4
P	Dascyllus albisella	10	KAHE7C	5	4.02		
P	Chromis vanderbilti	10	KAHE7C	4	1.26	9	5.28 0.2
	TOTAL	10	KAHE7C	46	2339.81	46	2339.81 100
C	Parupeneus multifasciatus	11	KAHE7D	1	11.05		
C	Parupeneus multifasciatus	11	KAHE7D	2	6.24		
C	Plectroglyphidodon johnstonianus	11	KAHE7D	1	0.86		
C	Paracirrhites arcatus	11	KAHE7D	1	3.45		
C	Paracirrhites arcatus	11	KAHE7D	1	8.12		
C	Cheilinus bimaculatus	11	KAHE7D	1	19.37		
C	Coris venusta	11	KAHE7D	1	1.01		
C	Stethojulis balteata	11	KAHE7D	1	4.00		
C	Sufflamen bursa	11	KAHE7D	1	85.87	10	139.97 36.8
CF	Chaetodon multicinctus	11	KAHE7D	1	0.83	1	0.83 0.2
H	Acanthurus nigrofuscus	11	KAHE7D	3	3.13		
H	Acanthurus blochii	11	KAHE7D	2	0.91		
H	Naso unicornis	11	KAHE7D	1	0.76	6	4.80 1.3
O	Melichthys vidua	11	KAHE7D	1	199.03		
O	Cantherhines sandwichiensis	11	KAHE7D	1	21.85		
O	Canthigaster jactator	11	KAHE7D	1	2.21		
O	Canthigaster jactator	11	KAHE7D	2	7.12	5	230.21 60.4
P	Chromis vanderbilti	11	KAHE7D	16	5.05	16	5.05 1.3
	TOTAL	11	KAHE7D	38	380.86	38	380.86 100
C	Saurida gracilis	12	KAHE7E	1	29.46		
C	Aulostomus chinensis	12	KAHE7E	1	140.29		
C	Decapterus macarellus	12	KAHE7E	150	40401.13		
C	Decapterus macarellus	12	KAHE7E	120	15633.49		
C	Parupeneus pleurostigma	12	KAHE7E	23	94.44		
C	Parupeneus multifasciatus	12	KAHE7E	10	110.52		
C	Parupeneus multifasciatus	12	KAHE7E	5	135.58		
C	Parupeneus multifasciatus	12	KAHE7E	3	163.20		
C	Parupeneus multifasciatus	12	KAHE7E	14	43.67		
C	Paracirrhites arcatus	12	KAHE7E	4	32.47		
C	Cirrhites fasciatus	12	KAHE7E	1	3.75		
C	Cheilinus bimaculatus	12	KAHE7E	1	4.10		
C	Thalassoma duperrey	12	KAHE7E	1	27.39		
C	Thalassoma duperrey	12	KAHE7E	4	219.80		
C	Thalassoma duperrey	12	KAHE7E	3	291.16		
C	Coris venusta	12	KAHE7E	9	22.99		
C	Pseudojuloides cerasinus	12	KAHE7E	1	6.32		
C	Pseudojuloides cerasinus	12	KAHE7E	1	1.28		
C	Macropharyngodon geoffroy	12	KAHE7E	3	7.49		
C	Zanclus cornutus	12	KAHE7E	1	104.16		
C	Sufflamen bursa	12	KAHE7E	6	515.24		
C	Sufflamen bursa	12	KAHE7E	3	2.58		
C	Sufflamen fraenatus	12	KAHE7E	1	224.79	366	58215.29 78.3
CF	Chaetodon multicinctus	12	KAHE7E	13	10.74		
CF	Chaetodon multicinctus	12	KAHE7E	4	26.63		
CF	Pervagor melanocephalus	12	KAHE7E	1	2.34	18	39.71 0.1
H	Scarus psittacus	12	KAHE7E	1	241.42		
H	Scarus psittacus	12	KAHE7E	10	378.73		
H	Scarus psittacus	12	KAHE7E	3	1664.07		
H	Scarus psittacus	12	KAHE7E	2	158.54		
H	Scarus psittacus	12	KAHE7E	2	1572.31		
H	Acanthurus nigrofuscus	12	KAHE7E	1	23.90		
H	Acanthurus nigrofuscus	12	KAHE7E	8	26.52		
H	Acanthurus nigrofuscus	12	KAHE7E	28	29.22		
H	Acanthurus olivaceus	12	KAHE7E	8	750.15		
H	Acanthurus olivaceus	12	KAHE7E	1	261.67		
H	Acanthurus olivaceus	12	KAHE7E	5	817.58		
H	Zebrasoma flavescens	12	KAHE7E	3	5.06		
H	Zebrasoma flavescens	12	KAHE7E	6	99.19		
H	Naso lituratus	12	KAHE7E	8	13.90		
H	Naso unicornis	12	KAHE7E	2	246.24		
H	Naso unicornis	12	KAHE7E	15	11.39	103	6299.87 8.5
O	Melichthys niger	12	KAHE7E	37	6043.41		
O	Melichthys vidua	12	KAHE7E	12	1492.15		
O	Melichthys vidua	12	KAHE7E	1	199.03		
O	Cantherhines sandwichiensis	12	KAHE7E	1	131.57		
O	Canthigaster jactator	12	KAHE7E	4	14.25	55	7880.41 10.6
P	Chaetodon kleini	12	KAHE7E	10	8.85		
P	Dascyllus albisella	12	KAHE7E	1	0.08		
P	Chromis vanderbilti	12	KAHE7E	209	66.02		
P	Naso brevirostris	12	KAHE7E	35	1362.36		
P	Naso brevirostris	12	KAHE7E	7	513.51	262	1950.81 2.6
	TOTAL	12	KAHE7E	804	74386.09	804	74386.09 100

20-Sep-13						GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS PERCENT
C	Aulostomus chinensis	13	KAHE10	1	19.27		
C	Aulostomus chinensis	13	KAHE10	1	8.10		
C	Aulostomus chinensis	13	KAHE10	2	421.24		
C	Apogon kallopterus	13	KAHE10	4	91.19		
C	Decapterus macarellus	13	KAHE10	23	4407.57		
C	Aphareus furcatus	13	KAHE10	1	5.74		
C	Monotaxis grandoculis	13	KAHE10	1	34.79		
C	Mulloides vanicolensis	13	KAHE10	65	3283.68		
C	Parupeneus pleurostigma	13	KAHE10	1	30.44		
C	Parupeneus multifasciatus	13	KAHE10	6	326.40		
C	Parupeneus multifasciatus	13	KAHE10	3	81.35		
C	Parupeneus multifasciatus	13	KAHE10	12	132.62		
C	Parupeneus multifasciatus	13	KAHE10	2	310.84		
C	Parupeneus bifasciatus	13	KAHE10	1	59.77		
C	Parupeneus cyclostomus	13	KAHE10	1	2.81		
C	Forcipiger flavissimus	13	KAHE10	3	7.59		
C	Forcipiger flavissimus	13	KAHE10	1	9.15		
C	Chaetodon fremblii	13	KAHE10	1	20.91		
C	Plectroglyphidodon johnstonianus	13	KAHE10	5	8.61		
C	Paracirrhites arcatus	13	KAHE10	6	48.71		
C	Paracirrhites forsteri	13	KAHE10	1	16.35		
C	Cirrhitis fasciatus	13	KAHE10	1	3.75		
C	Thalassoma duperrey	13	KAHE10	13	714.34		
C	Thalassoma duperrey	13	KAHE10	13	145.12		
C	Thalassoma duperrey	13	KAHE10	15	1455.78		
C	Thalassoma duperrey	13	KAHE10	18	493.03		
C	Gomphosus varius	13	KAHE10	3	12.07		
C	Gomphosus varius	13	KAHE10	1	22.60		
C	Stethojulis balteata	13	KAHE10	2	144.78		
C	Stethojulis balteata	13	KAHE10	8	115.27		
C	Stethojulis balteata	13	KAHE10	1	35.76		
C	Halichoeres ornatissimus	13	KAHE10	1	25.14		
C	Zanclus cornutus	13	KAHE10	4	96.17		
C	Sufflamen bursa	13	KAHE10	4	343.49	225	12934.41
CF	Exallias brevis	13	KAHE10	1	16.35	1	16.35
H	Calotomus carolinus	13	KAHE10	1	3318.76		53.6
H	Acanthurus triostegus	13	KAHE10	2	18.30		0.1
H	Acanthurus triostegus	13	KAHE10	6	25.43		
H	Acanthurus leucopareius	13	KAHE10	1	7.89		
H	Acanthurus nigrofusus	13	KAHE10	1	54.25		
H	Acanthurus nigrofusus	13	KAHE10	14	105.34		
H	Acanthurus nigrofusus	13	KAHE10	6	143.38		
H	Acanthurus nigrofusus	13	KAHE10	5	71.06		
H	Acanthurus olivaceus	13	KAHE10	8	2093.34		
H	Acanthurus olivaceus	13	KAHE10	6	562.61		
H	Acanthurus olivaceus	13	KAHE10	4	2252.68		
H	Acanthurus blochii	13	KAHE10	1	56.70		
H	Ctenochaetus strigosus	13	KAHE10	1	7.58		
H	Naso lituratus	13	KAHE10	8	13.90		
H	Naso unicornis	13	KAHE10	2	4.80	66	8736.03
O	Stegastes fasciolatus	13	KAHE10	17	250.97		36.2
O	Melichthys niger	13	KAHE10	2	326.67		
O	Melichthys vidua	13	KAHE10	1	199.03		
O	Canthigaster jactator	13	KAHE10	1	3.56	21	780.24
P	Dascyllus albisella	13	KAHE10	30	92.65		3.2
P	Abudefduf abdominalis	13	KAHE10	23	720.65		
P	Abudefduf vaigensis	13	KAHE10	23	720.65		
P	Chromis ovalis	13	KAHE10	6	35.59		
P	Chromis ovalis	13	KAHE10	32	80.31		
P	Chromis hanui	13	KAHE10	5	3.73	119	1653.58
	TOTAL	13	KAHE10	432	24120.62	432	24120.62
C	Parupeneus pleurostigma	14	NANA1	4	232.03		
C	Parupeneus pleurostigma	14	NANA1	3	39.76		
C	Parupeneus multifasciatus	14	NANA1	11	298.28		
C	Parupeneus multifasciatus	14	NANA1	5	15.60		
C	Parupeneus multifasciatus	14	NANA1	2	192.16		
C	Parupeneus multifasciatus	14	NANA1	8	435.20		
C	Parupeneus multifasciatus	14	NANA1	2	22.10		
C	Parupeneus bifasciatus	14	NANA1	1	11.60		
C	Parupeneus bifasciatus	14	NANA1	1	29.20		
C	Plectroglyphidodon imparipennis	14	NANA1	2	1.72		
C	Thalassoma duperrey	14	NANA1	10	273.90		
C	Thalassoma duperrey	14	NANA1	5	15.75		
C	Thalassoma duperrey	14	NANA1	5	55.82		
C	Thalassoma duperrey	14	NANA1	10	549.49		
C	Thalassoma lutescens	14	NANA1	1	32.76		
C	Stethojulis balteata	14	NANA1	4	143.05		
C	Stethojulis balteata	14	NANA1	8	115.27		
C	Rhinecanthus rectangulus	14	NANA1	1	45.36		
C	Rhinecanthus rectangulus	14	NANA1	2	171.75	85	2680.81
H	Acanthurus triostegus	14	NANA1	6	103.00		43.5
H	Acanthurus triostegus	14	NANA1	1	46.32		
H	Acanthurus triostegus	14	NANA1	2	8.48		
H	Acanthurus leucopareius	14	NANA1	6	94.16		
H	Acanthurus nigrofusus	14	NANA1	7	23.20		
H	Acanthurus nigrofusus	14	NANA1	4	56.85		
H	Acanthurus nigrofusus	14	NANA1	9	67.72		
H	Acanthurus olivaceus	14	NANA1	8	1308.13		
H	Acanthurus olivaceus	14	NANA1	1	5.73		
H	Acanthurus olivaceus	14	NANA1	4	45.29		
H	Acanthurus olivaceus	14	NANA1	3	1179.63		
H	Acanthurus olivaceus	14	NANA1	3	281.31		
H	Acanthurus blochii	14	NANA1	5	18.14		
H	Acanthurus blochii	14	NANA1	4	226.80		
H	Ctenochaetus strigosus	14	NANA1	3	9.26		
H	Naso unicornis	14	NANA1	1	2.40	67	3476.41
O	Canthigaster jactator	14	NANA1	2	7.12	2	7.12
	TOTAL	14	NANA1	154	6164.35	154	6164.35

20-Sep-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Parupeneus multifasciatus	15	NANA2	4	108.47				
C	Parupeneus multifasciatus	15	NANA2	1	96.08				
C	Parupeneus multifasciatus	15	NANA2	2	108.80				
C	Parupeneus bifasciatus	15	NANA2	1	59.77				
C	Chaetodon lunula	15	NANA2	3	107.98				
C	Plectroglyphidodon johnstonianus	15	NANA2	4	6.89				
C	Paracirrhites forsteri	15	NANA2	1	39.65				
C	Bodianus bilunulatus	15	NANA2	1	75.64				
C	Thalassoma duperrey	15	NANA2	9	873.47				
C	Thalassoma duperrey	15	NANA2	9	494.54				
C	Thalassoma duperrey	15	NANA2	5	136.95				
C	Coris venusta	15	NANA2	1	48.38				
C	Coris venusta	15	NANA2	1	23.64				
C	Stethojulis balteata	15	NANA2	3	43.23				
C	Stethojulis balteata	15	NANA2	2	144.78				
C	Halichoeres ornatissimus	15	NANA2	3	75.41				
C	Zanclus cornutus	15	NANA2	1	104.16				
C	Rhinecanthus rectangulus	15	NANA2	1	85.87				
C	Ostracion meleagris	15	NANA2	1	6.76	53	2640.46	10.1	
CF	Chaetodon ornatissimus	15	NANA2	1	0.98				
CF	Chaetodon ornatissimus	15	NANA2	2	138.00				
CF	Chaetodon quadrimaculatus	15	NANA2	2	50.61				
CF	Chaetodon multicinctus	15	NANA2	2	13.31				
CF	Exallias brevis	15	NANA2	1	3.45				
CF	Cantherhines dumerili	15	NANA2	1	117.96	9	324.31	1.2	
H	Calotomus carolinus	15	NANA2	1	499.89				
H	Acanthurus triostegus	15	NANA2	14	648.41				
H	Acanthurus leucopareus	15	NANA2	16	3722.80				
H	Acanthurus leucopareus	15	NANA2	5	333.70				
H	Acanthurus leucopareus	15	NANA2	42	5573.41				
H	Acanthurus leucopareus	15	NANA2	4	62.77				
H	Acanthurus nigrofuscus	15	NANA2	12	39.77				
H	Acanthurus nigrofuscus	15	NANA2	15	213.19				
H	Acanthurus nigrofuscus	15	NANA2	4	217.01				
H	Acanthurus olivaceus	15	NANA2	17	9573.89				
H	Ctenochaetus strigosus	15	NANA2	6	161.06				
H	Ctenochaetus strigosus	15	NANA2	18	55.58				
H	Ctenochaetus strigosus	15	NANA2	10	75.76				
H	Zebrasoma flavescens	15	NANA2	2	3.38				
H	Zebrasoma flavescens	15	NANA2	3	159.84				
H	Naso lituratus	15	NANA2	1	311.61	170	21652.10	82.6	
O	Stegastes fasciolatus	15	NANA2	3	44.29				
O	Melichthys niger	15	NANA2	6	980.01				
O	Cantherhines sandwichiensis	15	NANA2	1	82.05				
O	Canthigaster jactator	15	NANA2	1	3.56	11	1109.92	4.2	
P	Chromis vanderbilti	15	NANA2	12	3.79				
P	Naso hexacanthus	15	NANA2	4	492.47	16	496.26	1.9	
	TOTAL	15	NANA2	259	26223.05	259	26223.05	100	
C	Myripristis amaenus	16	PIPE	9	738.75				
C	Aulostomus chinensis	16	PIPE	1	27.72				
C	Decapterus macarellus	16	PIPE	45	8623.51				
C	Lutjanus kasmira	16	PIPE	18	1979.26				
C	Mulloides vanicolensis	16	PIPE	45	2273.32				
C	Parupeneus pleurostigma	16	PIPE	3	174.02				
C	Parupeneus multifasciatus	16	PIPE	3	288.24				
C	Parupeneus multifasciatus	16	PIPE	3	163.20				
C	Parupeneus cyclostomus	16	PIPE	1	10.05				
C	Forcipiger flavissimus	16	PIPE	1	9.15				
C	Paracirrhites arcatus	16	PIPE	3	49.05				
C	Thalassoma duperrey	16	PIPE	3	82.17				
C	Thalassoma duperrey	16	PIPE	21	2038.09				
C	Thalassoma duperrey	16	PIPE	12	659.39				
C	Gomphosus varius	16	PIPE	2	45.20				
C	Gomphosus varius	16	PIPE	1	62.03				
C	Stethojulis balteata	16	PIPE	1	72.39				
C	Stethojulis balteata	16	PIPE	4	143.05				
C	Halichoeres ornatissimus	16	PIPE	1	16.45				
C	Zanclus cornutus	16	PIPE	3	312.47				
C	Zanclus cornutus	16	PIPE	6	144.26				
C	Sufflamen bursa	16	PIPE	1	85.87	187	17997.64	21.8	
CF	Chaetodon multicinctus	16	PIPE	4	52.12				
CF	Pervagor melanocephalus	16	PIPE	1	9.75				
CF	Cantherhines dumerili	16	PIPE	1	194.98	6	256.85	0.3	
H	Calotomus carolinus	16	PIPE	1	1676.08				
H	Calotomus carolinus	16	PIPE	1	218.67				
H	Calotomus carolinus	16	PIPE	1	339.30				
H	Scarus psittacus	16	PIPE	34	4927.95				
H	Scarus psittacus	16	PIPE	27	10141.41				
H	Scarus psittacus	16	PIPE	19	4587.07				
H	Scarus psittacus	16	PIPE	4	3144.61				
H	Scarus rubroviolaceus	16	PIPE	1	1360.78				
H	Acanthurus triostegus	16	PIPE	5	231.58				
H	Acanthurus leucopareus	16	PIPE	3	398.10				
H	Acanthurus nigrofuscus	16	PIPE	26	621.32				
H	Acanthurus nigrofuscus	16	PIPE	18	976.54				
H	Acanthurus nigroris	16	PIPE	4	409.90				
H	Acanthurus olivaceus	16	PIPE	25	2344.22				
H	Acanthurus olivaceus	16	PIPE	22	5756.68				
H	Acanthurus dussumieri	16	PIPE	1	326.59				
H	Acanthurus blochii	16	PIPE	3	466.75	195	37927.55	46.0	
O	Stegastes fasciolatus	16	PIPE	23	597.55				
O	Canthigaster jactator	16	PIPE	7	24.93	30	622.48	0.8	
P	Chaetodon miliaris	16	PIPE	9	190.51				
P	Dascyllus albisella	16	PIPE	7	21.62				
P	Abudefduf abdominalis	16	PIPE	210	6579.84				
P	Abudefduf vaigensis	16	PIPE	425	13316.35				
P	Chromis ovalis	16	PIPE	485	5606.44	1136	25714.76	31.2	
	TOTAL	16	PIPE	1554	82519.27	1554	82519.27	100	

18 DECEMBER 2013 FIELD DATA

18-Dec-13

GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	GROUP BIOMASS	GROUP PERCENT
C	Parupeneus multifasciatus	1	EAST1	2	192.16			
C	Parupeneus multifasciatus	1	EAST1	1	27.12			
C	Plectroglyphidodon johnstonianus	1	EAST1	1	1.72			
C	Plectroglyphidodon imparipennis	1	EAST1	2	1.72			
C	Thalassoma duperrey	1	EAST1	12	659.39			
C	Thalassoma duperrey	1	EAST1	15	410.86			
C	Thalassoma duperrey	1	EAST1	7	679.36			
C	Thalassoma duperrey	1	EAST1	17	189.78			
C	Stethojulis balteata	1	EAST1	4	57.64			
C	Stethojulis balteata	1	EAST1	3	107.29			
C	Rhinecanthus rectangulus	1	EAST1	3	257.62			
C	Rhinecanthus rectangulus	1	EAST1	2	90.72			
C	Sufflamen fraenatus	1	EAST1	1	144.65			
C	Sufflamen fraenatus	1	EAST1	1	461.25	71	3281.28	14.3
CF	Chaetodon quadrimaculatus	1	EAST1	2	50.61	2	50.61	0.2
H	Calotomus carolinus	1	EAST1	1	1288.03			
H	Scarus rubroviolaceus	1	EAST1	2	2320.02			
H	Acanthurus triostegus	1	EAST1	15	1500.21			
H	Acanthurus triostegus	1	EAST1	23	1065.25			
H	Acanthurus leucopareius	1	EAST1	5	333.70			
H	Acanthurus nigrofusus	1	EAST1	23	1247.81			
H	Acanthurus nigrofusus	1	EAST1	18	430.15			
H	Acanthurus olivaceus	1	EAST1	7	3942.19			
H	Acanthurus dussumieri	1	EAST1	8	1904.70			
H	Acanthurus dussumieri	1	EAST1	9	634.90			
H	Acanthurus blochii	1	EAST1	21	2057.50			
H	Acanthurus blochii	1	EAST1	5	2267.97			
H	Naso lituratus	1	EAST1	2	255.47	139	19247.89	83.9
O	Melichthys vidua	1	EAST1	1	296.24			
O	Canthigaster jactator	1	EAST1	3	10.69	4	306.92	1.3
P	Chaetodon miliaris	1	EAST1	1	21.17			
P	Chromis vanderbilti	1	EAST1	13	4.11			
P	Naso brevirostris	1	EAST1	1	38.92	15	64.20	0.3
	TOTAL	1	EAST1	231	22950.91	231	22950.91	100
C	Lutjanus fulvus	2	EAST2	1	155.61			
C	Parupeneus pleurostigma	2	EAST2	1	58.01			
C	Parupeneus multifasciatus	2	EAST2	1	340.44			
C	Parupeneus multifasciatus	2	EAST2	1	27.12			
C	Parupeneus bifasciatus	2	EAST2	1	59.77			
C	Parupeneus bifasciatus	2	EAST2	1	29.20			
C	Parupeneus cyclostomus	2	EAST2	1	316.39			
C	Paracirrhites arcatus	2	EAST2	1	16.35			
C	Thalassoma duperrey	2	EAST2	13	356.08			
C	Thalassoma duperrey	2	EAST2	11	1067.57			
C	Thalassoma duperrey	2	EAST2	15	824.23			
C	Gomphosus varius	2	EAST2	1	39.39			
C	Gomphosus varius	2	EAST2	1	11.04			
C	Stethojulis balteata	2	EAST2	3	107.29			
C	Sufflamen bursa	2	EAST2	1	85.87	53	3494.36	6.2
CF	Chaetodon unimaculatus	2	EAST2	4	101.22			
CF	Chaetodon ornatissimus	2	EAST2	2	138.00			
CF	Chaetodon quadrimaculatus	2	EAST2	6	151.82	12	391.05	0.7
H	Scarus sordidus	2	EAST2	5	68.65			
H	Scarus rubroviolaceus	2	EAST2	5	1732.52			
H	Scarus rubroviolaceus	2	EAST2	6	4042.52			
H	Scarus rubroviolaceus	2	EAST2	6	6960.05			
H	Acanthurus triostegus	2	EAST2	3	300.04			
H	Acanthurus triostegus	2	EAST2	12	555.78			
H	Acanthurus leucopareius	2	EAST2	2	265.40			
H	Acanthurus nigrofusus	2	EAST2	28	210.69			
H	Acanthurus nigrofusus	2	EAST2	22	525.73			
H	Acanthurus olivaceus	2	EAST2	38	21400.47			
H	Acanthurus olivaceus	2	EAST2	61	15961.69			
H	Ctenochaetus strigosus	2	EAST2	3	80.53			
H	Zebrasoma flavescens	2	EAST2	1	53.28			
H	Naso lituratus	2	EAST2	1	127.73	193	52285.10	93.1
O	Canthigaster jactator	2	EAST2	3	6.62	3	6.62	0.01
	TOTAL	2	EAST2	261	56177.12	261	56177.12	100

18-Dec-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Decapterus macarellus	3	EAST3	13	1693.63				
C	Parupeneus multifasciatus	3	EAST3	1	27.12				
C	Parupeneus multifasciatus	3	EAST3	1	96.08				
C	Parupeneus bifasciatus	3	EAST3	2	119.53				
C	Plectroglyphidodon johnstonianus	3	EAST3	5	8.61				
C	Paracirrhites arcatus	3	EAST3	1	16.35				
C	Paracirrhites forsteri	3	EAST3	1	39.65				
C	Labroides phthiropagus	3	EAST3	2	1.26				
C	Pseudochelinus octotaenia	3	EAST3	2	71.53				
C	Thalassoma duperrey	3	EAST3	2	194.10				
C	Thalassoma duperrey	3	EAST3	8	439.59				
C	Thalassoma duperrey	3	EAST3	6	164.34				
C	Gomphosus varius	3	EAST3	1	22.60				
C	Stethojulis balteata	3	EAST3	3	107.29				
C	Sufflamen bursa	3	EAST3	1	85.87				
C	Sufflamen fraenatus	3	EAST3	1	144.65	50	3232.20	25.3	
CF	Chaetodon unimaculatus	3	EAST3	3	42.67				
CF	Chaetodon ornatissimus	3	EAST3	2	138.00				
CF	Chaetodon multicinctus	3	EAST3	6	78.18				
CF	Chaetodon multicinctus	3	EAST3	1	0.83	12	259.68	2.0	
H	Calotomus carolinus	3	EAST3	2	69.38				
H	Scarus psittacus	3	EAST3	4	58.46				
H	Scarus psittacus	3	EAST3	1	786.15				
H	Scarus rubroviolaceus	3	EAST3	1	1472.49				
H	Acanthurus nigrofuscus	3	EAST3	13	184.77				
H	Acanthurus nigrofuscus	3	EAST3	17	406.25				
H	Acanthurus nigrofuscus	3	EAST3	10	75.25				
H	Acanthurus olivaceus	3	EAST3	4	2252.68				
H	Acanthurus olivaceus	3	EAST3	12	3140.00				
H	Ctenochaetus strigosus	3	EAST3	16	243.18				
H	Ctenochaetus strigosus	3	EAST3	8	214.75				
H	Ctenochaetus strigosus	3	EAST3	4	30.30				
H	Zebrasoma flavescens	3	EAST3	1	9.48				
H	Zebrasoma flavescens	3	EAST3	1	1.69				
H	Zebrasoma flavescens	3	EAST3	2	106.56	96	9051.39	71.0	
O	Stegastes fasciolatus	3	EAST3	6	88.58				
O	Cantherhines sandwichiensis	3	EAST3	1	82.05	7	170.63	1.3	
P	Chaetodon miliaris	3	EAST3	2	42.34	2	42.34	0.3	
	TOTAL	3	EAST3	167	12756.23	167	12756.23	100	
C	Scorpaenopsis diabolus	4	EAST4	1	2684.85				
C	Parupeneus multifasciatus	4	EAST4	1	27.12				
C	Parupeneus multifasciatus	4	EAST4	1	96.08				
C	Plectroglyphidodon johnstonianus	4	EAST4	2	3.44				
C	Plectroglyphidodon imparipennis	4	EAST4	1	0.86				
C	Paracirrhites arcatus	4	EAST4	4	65.39				
C	Thalassoma duperrey	4	EAST4	15	824.23				
C	Thalassoma duperrey	4	EAST4	12	1164.62				
C	Thalassoma duperrey	4	EAST4	21	575.20				
C	Stethojulis balteata	4	EAST4	1	35.76				
C	Stethojulis balteata	4	EAST4	1	72.39				
C	Halichoeres ornatissimus	4	EAST4	3	28.57				
C	Halichoeres ornatissimus	4	EAST4	3	75.41				
C	Rhinecanthus rectangulus	4	EAST4	1	85.87				
C	Rhinecanthus rectangulus	4	EAST4	1	45.36				
C	Sufflamen fraenatus	4	EAST4	1	461.25	69	6246.43	23.2	
CF	Chaetodon quadrimaculatus	4	EAST4	2	50.61	2	50.61	0.2	
H	Acanthurus nigrofuscus	4	EAST4	37	525.87				
H	Acanthurus nigrofuscus	4	EAST4	16	120.39				
H	Acanthurus nigrofuscus	4	EAST4	28	669.12				
H	Acanthurus olivaceus	4	EAST4	19	4971.67				
H	Acanthurus olivaceus	4	EAST4	18	10137.06				
H	Acanthurus blochii	4	EAST4	1	453.59				
H	Ctenochaetus strigosus	4	EAST4	1	15.20	120	16892.91	62.7	
O	Stegastes fasciolatus	4	EAST4	2	51.96				
O	Melichthys niger	4	EAST4	12	1960.03				
O	Melichthys vidua	4	EAST4	3	1699.63				
O	Canthigaster jactator	4	EAST4	2	7.12	19	3718.74	13.8	
P	Chaetodon miliaris	4	EAST4	1	21.17				
P	Chromis vanderbilti	4	EAST4	40	12.64	41	33.80	0.1	
	TOTAL	4	EAST4	251	26942.50	251	26942.50	100	

18-Dec-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Parupeneus multifasciatus	5	KO1	1	155.42				
C	Parupeneus multifasciatus	5	KO1	1	54.40				
C	Plectroglyphidodon johnstonianus	5	KO1	2	3.44				
C	Paracirrhites arcatus	5	KO1	3	24.35				
C	Cheilinus rhodochrous	5	KO1	1	30.14				
C	Thalassoma duperrey	5	KO1	2	194.10				
C	Thalassoma duperrey	5	KO1	6	164.34				
C	Thalassoma duperrey	5	KO1	4	44.65				
C	Thalassoma duperrey	5	KO1	2	109.90				
C	Thalassoma ballieui	5	KO1	1	167.53				
C	Gomphosus varius	5	KO1	1	22.60				
C	Stethojulis balteata	5	KO1	4	143.05				
C	Stethojulis balteata	5	KO1	3	217.17				
C	Halichoeres ornatissimus	5	KO1	2	50.27				
C	Halichoeres ornatissimus	5	KO1	1	16.45				
C	Sufflamen bursa	5	KO1	1	144.65				
C	Sufflamen bursa	5	KO1	3	257.62	38	1800.10	23.1	
CF	Chaetodon unimaculatus	5	KO1	2	50.61				
CF	Chaetodon multicinctus	5	KO1	4	52.12	6	102.73	1.3	
H	Scarus sordidus	5	KO1	1	367.02				
H	Acanthurus triostegus	5	KO1	6	103.00				
H	Acanthurus leucopareius	5	KO1	1	232.68				
H	Acanthurus leucopareius	5	KO1	1	132.70				
H	Acanthurus nigrofusus	5	KO1	13	310.66				
H	Acanthurus nigrofusus	5	KO1	15	813.79				
H	Acanthurus olivaceus	5	KO1	3	785.00				
H	Ctenochaetus strigosus	5	KO1	22	1449.03				
H	Ctenochaetus strigosus	5	KO1	29	778.47				
H	Ctenochaetus strigosus	5	KO1	13	197.58	104	5169.93	66.3	
O	Stegastes fasciolatus	5	KO1	8	207.84				
O	Melichthys niger	5	KO1	2	496.06				
O	Canthigaster jactator	5	KO1	7	24.93	17	728.84	9.3	
	TOTAL	5	KO1	165	7801.59	165	7801.59	100	
C	Aulostomus chinensis	6	KO2	1	52.00				
C	Parupeneus multifasciatus	6	KO2	1	96.08				
C	Parupeneus multifasciatus	6	KO2	1	27.12				
C	Parupeneus cyclostomus	6	KO2	1	440.45				
C	Parupeneus cyclostomus	6	KO2	1	780.78				
C	Forcipiger flavissimus	6	KO2	2	18.30				
C	Plectroglyphidodon johnstonianus	6	KO2	1	1.72				
C	Labroides phthiophagus	6	KO2	2	1.26				
C	Thalassoma duperrey	6	KO2	5	274.74				
C	Thalassoma duperrey	6	KO2	5	136.95				
C	Thalassoma duperrey	6	KO2	2	194.10				
C	Gomphosus varius	6	KO2	2	45.20				
C	Stethojulis balteata	6	KO2	1	72.39				
C	Macropharyngodon geoffroy	6	KO2	2	37.26				
C	Sufflamen bursa	6	KO2	2	171.75				
C	Sufflamen bursa	6	KO2	1	144.65	30	2494.75	11.4	
CF	Chaetodon unimaculatus	6	KO2	6	151.82				
CF	Chaetodon quadrimaculatus	6	KO2	4	101.22				
CF	Chaetodon multicinctus	6	KO2	2	13.31	12	266.35	1.2	
H	Scarus psittacus	6	KO2	6	87.69				
H	Scarus psittacus	6	KO2	8	302.98				
H	Scarus rubroviolaceus	6	KO2	1	43.92				
H	Scarus rubroviolaceus	6	KO2	2	2944.98				
H	Acanthurus triostegus	6	KO2	5	85.83				
H	Acanthurus triostegus	6	KO2	9	416.84				
H	Acanthurus leucopareius	6	KO2	1	132.70				
H	Acanthurus leucopareius	6	KO2	2	133.48				
H	Acanthurus nigrofusus	6	KO2	15	213.19				
H	Acanthurus nigrofusus	6	KO2	16	868.04				
H	Acanthurus nigrofusus	6	KO2	12	286.76				
H	Acanthurus nigroris	6	KO2	3	162.76				
H	Acanthurus nigroris	6	KO2	2	204.95				
H	Acanthurus olivaceus	6	KO2	7	656.38				
H	Acanthurus olivaceus	6	KO2	2	1126.34				
H	Acanthurus olivaceus	6	KO2	22	5756.68				
H	Acanthurus blochii	6	KO2	3	87.09				
H	Ctenochaetus strigosus	6	KO2	11	724.51				
H	Ctenochaetus strigosus	6	KO2	10	1321.34				
H	Ctenochaetus strigosus	6	KO2	10	268.44				
H	Zebrasoma flavescens	6	KO2	5	266.40				
H	Zebrasoma flavescens	6	KO2	17	161.23				
H	Zebrasoma flavescens	6	KO2	8	208.24	177	16460.76	75.5	
O	Stegastes fasciolatus	6	KO2	3	44.29				
O	Melichthys niger	6	KO2	5	816.68				
O	Melichthys niger	6	KO2	2	496.06				
O	Melichthys vidua	6	KO2	1	199.03				
O	Canthigaster jactator	6	KO2	3	10.69	14	1566.75	7.2	
P	Abudefduf abdominalis	6	KO2	32	1002.64				
P	Chromis vanderbilti	6	KO2	13	4.11				
P	Chromis hanui	6	KO2	3	2.24	48	1008.99	4.6	
	TOTAL	6	KO2	281	21797.60	281	21797.60	99.9999998	

18-Dec-13							GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	GROUP PERCENT
C	Mulloides flavolineatus	7	KAHE1D	5	1826.04			
C	Parupeneus multifasciatus	7	KAHE1D	1	54.40			
C	Parupeneus multifasciatus	7	KAHE1D	1	27.12			
C	Parupeneus bifasciatus	7	KAHE1D	1	107.31			
C	Plectroglyphidodon imparipennis	7	KAHE1D	1	0.86			
C	Paracirrhites arcatus	7	KAHE1D	1	8.12			
C	Thalassoma duperrey	7	KAHE1D	8	219.12			
C	Thalassoma duperrey	7	KAHE1D	23	1263.82			
C	Thalassoma duperrey	7	KAHE1D	24	2329.25			
C	Gomphosus varius	7	KAHE1D	2	45.20			
C	Sufflamen bursa	7	KAHE1D	4	181.45			
C	Sufflamen bursa	7	KAHE1D	4	343.49	75	6406.19	31.3
CF	Chaetodon quadrimaculatus	7	KAHE1D	1	7.03			
CF	Cantherhines dumerili	7	KAHE1D	1	301.33	2	308.35	1.5
H	Scarus psittacus	7	KAHE1D	12	454.47			
H	Scarus psittacus	7	KAHE1D	16	233.83			
H	Acanthurus triostegus	7	KAHE1D	12	555.78			
H	Acanthurus triostegus	7	KAHE1D	35	600.83			
H	Acanthurus leucopareius	7	KAHE1D	1	27.52			
H	Acanthurus nigrofuscus	7	KAHE1D	39	931.98			
H	Acanthurus nigrofuscus	7	KAHE1D	10	142.13			
H	Ctenochaetus strigosus	7	KAHE1D	4	263.46			
H	Ctenochaetus strigosus	7	KAHE1D	7	106.39			
H	Naso unicornis	7	KAHE1D	1	5.44	137	3321.82	16.2
O	Stegastes fasciolatus	7	KAHE1D	2	51.96			
O	Melichthys niger	7	KAHE1D	63	10290.14			
O	Cantherhines sandwichiensis	7	KAHE1D	1	82.05			
O	Canthigaster jactator	7	KAHE1D	5	17.81	71	10441.96	50.9
P	Chromis vanderbilti	7	KAHE1D	52	16.43	52	16.43	0.1
	TOTAL	7	KAHE1D	337	20494.75	337	20494.75	100
C	Parupeneus pleurostigma	8	KAHE5B	1	58.01			
C	Parupeneus multifasciatus	8	KAHE5B	1	155.42			
C	Parupeneus multifasciatus	8	KAHE5B	3	81.35			
C	Parupeneus multifasciatus	8	KAHE5B	2	108.80			
C	Plectroglyphidodon johnstonianus	8	KAHE5B	3	5.17			
C	Paracirrhites arcatus	8	KAHE5B	6	98.09			
C	Novaculichthys taeniourus	8	KAHE5B	1	12.62			
C	Thalassoma duperrey	8	KAHE5B	5	55.82			
C	Thalassoma duperrey	8	KAHE5B	8	219.12			
C	Thalassoma duperrey	8	KAHE5B	8	439.59			
C	Thalassoma duperrey	8	KAHE5B	3	291.16			
C	Stethojulis balteata	8	KAHE5B	1	72.39			
C	Macropharyngodon geoffroy	8	KAHE5B	3	55.89			
C	Halichoeres ornatissimus	8	KAHE5B	2	32.90			
C	Rhinecanthus rectangulus	8	KAHE5B	1	45.36			
C	Sufflamen bursa	8	KAHE5B	6	515.24			
C	Ostracion meleagris	8	KAHE5B	1	6.76	55	2253.69	30.4
CF	Chaetodon unimaculatus	8	KAHE5B	2	28.45			
CF	Chaetodon multicinctus	8	KAHE5B	2	26.06			
CF	Chaetodon multicinctus	8	KAHE5B	2	13.31	6	67.82	0.9
H	Scarus psittacus	8	KAHE5B	14	530.22			
H	Acanthurus nigrofuscus	8	KAHE5B	25	597.43			
H	Acanthurus nigrofuscus	8	KAHE5B	16	227.40			
H	Acanthurus nigrofuscus	8	KAHE5B	2	108.50			
H	Acanthurus olivaceus	8	KAHE5B	4	2252.68			
H	Acanthurus olivaceus	8	KAHE5B	4	375.08			
H	Acanthurus olivaceus	8	KAHE5B	2	523.33			
H	Naso lituratus	8	KAHE5B	2	29.80			
H	Naso unicornis	8	KAHE5B	3	51.58			
H	Naso unicornis	8	KAHE5B	1	5.44	73	4701.46	63.5
O	Stegastes fasciolatus	8	KAHE5B	3	77.94			
O	Melichthys vidua	8	KAHE5B	1	199.03			
O	Cantherhines sandwichiensis	8	KAHE5B	1	82.05			
O	Canthigaster jactator	8	KAHE5B	4	14.25	9	373.27	5.0
P	Chromis vanderbilti	8	KAHE5B	20	6.32	20	6.32	0.1
	TOTAL	8	KAHE5B	163	7402.55	163	7402.55	100.000001
C	Myripristis amaenus	9	KAHE7B	1	28.56			
C	Scorpaenopsis diabolus	9	KAHE7B	1	341.72			
C	Parupeneus multifasciatus	9	KAHE7B	4	108.47			
C	Parupeneus multifasciatus	9	KAHE7B	3	163.20			
C	Parupeneus multifasciatus	9	KAHE7B	1	96.08			
C	Paracirrhites arcatus	9	KAHE7B	1	3.45			
C	Bodianus bilunulatus	9	KAHE7B	1	133.12			
C	Thalassoma duperrey	9	KAHE7B	1	11.16			
C	Thalassoma duperrey	9	KAHE7B	4	109.56			
C	Gomphosus varius	9	KAHE7B	1	62.03			
C	Halichoeres ornatissimus	9	KAHE7B	1	4.41			
C	Zanclus cornutus	9	KAHE7B	1	54.90			
C	Sufflamen bursa	9	KAHE7B	3	257.62	23	1374.26	53.7
CF	Pervagor melanocephalus	9	KAHE7B	1	5.23			
CF	Cantherhines dumerili	9	KAHE7B	1	194.98	2	200.21	7.8
H	Acanthurus triostegus	9	KAHE7B	1	17.17			
H	Acanthurus triostegus	9	KAHE7B	1	46.32			
H	Acanthurus nigrofuscus	9	KAHE7B	7	167.28			
H	Acanthurus nigrofuscus	9	KAHE7B	9	67.72			
H	Acanthurus olivaceus	9	KAHE7B	1	163.52			
H	Acanthurus olivaceus	9	KAHE7B	2	94.95			
H	Naso lituratus	9	KAHE7B	2	29.80			
H	Naso lituratus	9	KAHE7B	1	36.34			
H	Naso lituratus	9	KAHE7B	3	12.72			
H	Naso unicornis	9	KAHE7B	2	34.39			
H	Naso unicornis	9	KAHE7B	3	16.31	32	686.51	26.8
O	Melichthys vidua	9	KAHE7B	1	199.03			
O	Canthigaster jactator	9	KAHE7B	4	14.25	5	213.28	8.3
P	Dascyllus albisella	9	KAHE7B	1	8.03			
P	Chromis vanderbilti	9	KAHE7B	13	4.11			
P	Chromis hamui	9	KAHE7B	2	1.49			
P	Naso brevirostris	9	KAHE7B	1	73.36	17	86.98	3.4
	TOTAL	9	KAHE7B	79	2561.24	79	2561.24	100

18-Dec-13							GROUP	GROUP
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT
C	Fistularia commersoni	10	KAHE7C	1	14.94			
C	Decapterus macarellus	10	KAHE7C	8	1533.07			
C	Parupeneus pleurostigma	10	KAHE7C	6	182.63			
C	Parupeneus multifasciatus	10	KAHE7C	2	54.23			
C	Parupeneus multifasciatus	10	KAHE7C	3	163.20			
C	Parupeneus multifasciatus	10	KAHE7C	1	155.42			
C	Paracirrhites arcatus	10	KAHE7C	2	16.24			
C	Zanclus cornutus	10	KAHE7C	1	24.04			
C	Sufflamen bursa	10	KAHE7C	6	515.24	30	2659.01	70.3
CF	Chaetodon multicinctus	10	KAHE7C	2	1.65			
CF	Pervagor melanocephalus	10	KAHE7C	2	32.43	4	34.09	0.9
H	Acanthurus olivaceus	10	KAHE7C	1	393.21			
H	Naso lituratus	10	KAHE7C	6	25.43			
H	Naso lituratus	10	KAHE7C	6	89.39			
H	Naso unicornis	10	KAHE7C	9	154.75			
H	Naso unicornis	10	KAHE7C	4	21.74	26	684.53	18.1
O	Melichthys vidua	10	KAHE7C	1	199.03			
O	Cantherhines sandwichiensis	10	KAHE7C	1	82.05	2	281.08	7.4
P	Dascyllus albisella	10	KAHE7C	1	0.80			
P	Dascyllus albisella	10	KAHE7C	1	3.09			
P	Chromis vanderbilti	10	KAHE7C	3	0.95			
P	Naso brevirostris	10	KAHE7C	4	106.56			
P	Naso brevirostris	10	KAHE7C	2	10.87	11	122.27	3.2
	TOTAL	10	KAHE7C	73	3780.98	73	3780.98	100
C	Parupeneus multifasciatus	11	KAHE7D	1	54.40			
C	Parupeneus multifasciatus	11	KAHE7D	17	460.98			
C	Plectroglyphidodon johnstonianus	11	KAHE7D	1	1.72			
C	Paracirrhites arcatus	11	KAHE7D	2	16.24			
C	Paracirrhites arcatus	11	KAHE7D	1	16.35			
C	Thalassoma duperrey	11	KAHE7D	1	54.95	23	604.64	46.3
CF	Chaetodon multicinctus	11	KAHE7D	1	0.83	1	0.83	0.1
H	Calotomus carolinus	11	KAHE7D	1	72.28			
H	Calotomus carolinus	11	KAHE7D	1	13.46			
H	Acanthurus nigrofuscus	11	KAHE7D	1	7.52			
H	Acanthurus olivaceus	11	KAHE7D	6	34.39			
H	Acanthurus olivaceus	11	KAHE7D	8	157.95			
H	Naso lituratus	11	KAHE7D	1	4.24			
H	Naso unicornis	11	KAHE7D	4	21.74	22	311.60	23.9
O	Melichthys vidua	11	KAHE7D	1	199.03			
O	Canthigaster coronata	11	KAHE7D	1	22.05			
O	Canthigaster jactator	11	KAHE7D	2	7.12	4	228.20	17.5
P	Chaetodon kleini	11	KAHE7D	3	8.57			
P	Chromis vanderbilti	11	KAHE7D	4	1.26			
P	Naso brevirostris	11	KAHE7D	4	68.78			
P	Naso brevirostris	11	KAHE7D	15	81.54	26	160.15	12.3
	TOTAL	11	KAHE7D	76	1305.41	76	1305.41	100
C	Gymnothorax meleagris	12	KAHE7E	1	166.27			
C	Aulostomus chinensis	12	KAHE7E	1	52.00			
C	Parupeneus pleurostigma	12	KAHE7E	1	13.25			
C	Parupeneus pleurostigma	12	KAHE7E	9	273.94			
C	Parupeneus multifasciatus	12	KAHE7E	9	99.46			
C	Parupeneus multifasciatus	12	KAHE7E	7	189.82			
C	Parupeneus multifasciatus	12	KAHE7E	2	108.80			
C	Forcipiger flavissimus	12	KAHE7E	2	18.30			
C	Plectroglyphidodon johnstonianus	12	KAHE7E	2	1.72			
C	Paracirrhites arcatus	12	KAHE7E	2	6.89			
C	Paracirrhites forsteri	12	KAHE7E	1	39.65			
C	Cirrhitops fasciatus	12	KAHE7E	1	15.63			
C	Pseudocheilinus octotaenia	12	KAHE7E	1	35.76			
C	Thalassoma duperrey	12	KAHE7E	3	82.17			
C	Thalassoma duperrey	12	KAHE7E	2	22.33			
C	Thalassoma duperrey	12	KAHE7E	1	97.05			
C	Thalassoma quinquevittatum	12	KAHE7E	1	63.55			
C	Pseudojuloides cerasinus	12	KAHE7E	4	44.65			
C	Pseudojuloides cerasinus	12	KAHE7E	3	9.45			
C	Stethojulis balteata	12	KAHE7E	1	35.76			
C	Stethojulis balteata	12	KAHE7E	1	72.39			
C	Macropharyngodon geoffroy	12	KAHE7E	1	42.90			
C	Anampses chrysocephalus	12	KAHE7E	1	2.63			
C	Zanclus cornutus	12	KAHE7E	1	54.90			
C	Sufflamen bursa	12	KAHE7E	2	39.85			
C	Sufflamen bursa	12	KAHE7E	1	144.65			
C	Sufflamen bursa	12	KAHE7E	7	601.11			
C	Sufflamen fraenatus	12	KAHE7E	1	329.34			
C	Sufflamen fraenatus	12	KAHE7E	2	289.30	71	2953.54	54.9
CF	Chaetodon ornatissimus	12	KAHE7E	1	3.40			
CF	Chaetodon multicinctus	12	KAHE7E	8	22.40	9	25.80	0.5
H	Centropyge potteri	12	KAHE7E	1	1.72			
H	Acanthurus nigrofuscus	12	KAHE7E	20	150.49			
H	Acanthurus nigrofuscus	12	KAHE7E	33	109.38			
H	Acanthurus olivaceus	12	KAHE7E	2	786.42			
H	Acanthurus olivaceus	12	KAHE7E	1	19.74			
H	Ctenochaetus strigosus	12	KAHE7E	9	7.84			
H	Ctenochaetus strigosus	12	KAHE7E	3	9.26			
H	Zebrasoma flavescens	12	KAHE7E	1	1.69			
H	Naso lituratus	12	KAHE7E	2	29.80			
H	Naso lituratus	12	KAHE7E	5	21.19			
H	Naso unicornis	12	KAHE7E	1	5.44	78	1142.97	21.2
O	Melichthys niger	12	KAHE7E	3	490.01			
O	Melichthys vidua	12	KAHE7E	3	597.09			
O	Canthigaster jactator	12	KAHE7E	2	4.41	8	1091.51	20.3
P	Chaetodon kleini	12	KAHE7E	3	8.57			
P	Dascyllus albisella	12	KAHE7E	6	4.82			
P	Chromis vanderbilti	12	KAHE7E	171	54.02			
P	Chromis hanui	12	KAHE7E	22	16.43			
P	Naso brevirostris	12	KAHE7E	3	16.31			
P	Naso brevirostris	12	KAHE7E	4	68.78	209	168.92	3.1
	TOTAL	12	KAHE7E	375	5382.73	375	5382.73	100.000001

18-Dec-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Adioryx xantherythrus	13	KAHE10	1	31.67				
C	Aulostomus chinensis	13	KAHE10	1	210.62				
C	Aulostomus chinensis	13	KAHE10	1	4.76				
C	Fistularia commersoni	13	KAHE10	1	253.56				
C	Priacanthus meeki	13	KAHE10	1	250.49				
C	Mulloides flavolineatus	13	KAHE10	183	15270.37				
C	Parupeneus pleurostigma	13	KAHE10	7	406.06				
C	Parupeneus multifasciatus	13	KAHE10	2	54.23				
C	Parupeneus multifasciatus	13	KAHE10	1	96.08				
C	Parupeneus multifasciatus	13	KAHE10	2	108.80				
C	Parupeneus bifasciatus	13	KAHE10	1	107.31				
C	Parupeneus bifasciatus	13	KAHE10	2	119.53				
C	Forcipiger flavissimus	13	KAHE10	1	5.13				
C	Plectroglyphidodon johnstonianus	13	KAHE10	3	5.17				
C	Plectroglyphidodon imparipennis	13	KAHE10	1	0.86				
C	Paracirrhites arcatus	13	KAHE10	1	16.35				
C	Paracirrhites forsteri	13	KAHE10	1	16.35				
C	Cirrihitops fasciatus	13	KAHE10	1	8.23				
C	Thalassoma duperrey	13	KAHE10	2	54.78				
C	Thalassoma duperrey	13	KAHE10	5	485.26				
C	Thalassoma duperrey	13	KAHE10	9	494.54				
C	Thalassoma purpureum	13	KAHE10	1	165.99				
C	Coris gaimard	13	KAHE10	1	45.99				
C	Stethojulis balteata	13	KAHE10	1	72.39				
C	Stethojulis balteata	13	KAHE10	1	35.76				
C	Macropharyngodon geoffroy	13	KAHE10	1	29.13				
C	Halichoeres ornatissimus	13	KAHE10	2	50.27				
C	Zanclus cornutus	13	KAHE10	1	54.90				
C	Sufflamen bursa	13	KAHE10	1	19.92				
C	Sufflamen bursa	13	KAHE10	2	171.75				
C	Sufflamen fraenatus	13	KAHE10	1	224.79	239	18871.04	88.1	
CF	Chaetodon multicinctus	13	KAHE10	2	26.06				
CF	Pervagor spilosoma	13	KAHE10	1	16.22	3	42.28	0.2	
H	Acanthurus triostegus	13	KAHE10	3	51.50				
H	Acanthurus triostegus	13	KAHE10	1	4.24				
H	Acanthurus leucopareius	13	KAHE10	1	66.74				
H	Acanthurus nigrofuscus	13	KAHE10	5	37.62				
H	Acanthurus nigrofuscus	13	KAHE10	2	108.50				
H	Acanthurus nigrofuscus	13	KAHE10	7	167.28				
H	Acanthurus olivaceus	13	KAHE10	2	327.03				
H	Acanthurus olivaceus	13	KAHE10	3	785.00				
H	Acanthurus olivaceus	13	KAHE10	1	47.48				
H	Naso lituratus	13	KAHE10	2	16.93				
H	Naso lituratus	13	KAHE10	2	29.80				
H	Naso lituratus	13	KAHE10	1	36.34				
H	Naso unicornis	13	KAHE10	1	17.19	31	1695.66	7.9	
O	Stegastes fasciolatus	13	KAHE10	8	207.84				
O	Melichthys niger	13	KAHE10	2	326.67				
O	Cantherhines sandwichiensis	13	KAHE10	1	82.05	11	616.56	2.9	
P	Dascyllus albisella	13	KAHE10	12	37.06				
P	Chromis vanderbilti	13	KAHE10	28	8.84				
P	Chromis ovalis	13	KAHE10	12	138.72	52	184.62	0.9	
	TOTAL	13	KAHE10	336	21410.16	336	21410.16	99.9999998	
C	Gymnothorax meleagris	14	NANA1	1	24.76				
C	Parupeneus multifasciatus	14	NANA1	1	54.40				
C	Parupeneus multifasciatus	14	NANA1	1	27.12				
C	Parupeneus multifasciatus	14	NANA1	2	192.16				
C	Plectroglyphidodon imparipennis	14	NANA1	1	0.86				
C	Thalassoma duperrey	14	NANA1	3	291.16				
C	Thalassoma duperrey	14	NANA1	2	54.78				
C	Thalassoma duperrey	14	NANA1	4	219.80				
C	Thalassoma purpureum	14	NANA1	1	100.89				
C	Thalassoma fuscum	14	NANA1	1	63.55				
C	Stethojulis balteata	14	NANA1	1	35.76				
C	Rhinecanthus rectangulus	14	NANA1	1	45.36	19	1110.60	58.4	
H	Acanthurus triostegus	14	NANA1	3	138.95				
H	Acanthurus nigrofuscus	14	NANA1	2	108.50				
H	Acanthurus nigrofuscus	14	NANA1	14	105.34				
H	Acanthurus nigrofuscus	14	NANA1	17	406.25				
H	Acanthurus blochii	14	NANA1	1	12.25	37	771.29	40.5	
O	Stegastes fasciolatus	14	NANA1	1	14.76				
O	Canthigaster jactator	14	NANA1	1	3.56	2	18.33	1.0	
P	Chromis vanderbilti	14	NANA1	6	1.90	6	1.90	0.1	
	TOTAL	14	NANA1	64	1902.11	64	1902.11	100	

18-Dec-13						GROUP		GROUP	
GRP	SPECIES	RB	STATION	NO.	BIOMASS	NO. INDIV.	BIOMASS	PERCENT	
C	Fistularia commersoni	15	NANA2	1	100.51				
C	Parupeneus pleurostigma	15	NANA2	1	58.01				
C	Parupeneus multifasciatus	15	NANA2	2	54.23				
C	Parupeneus multifasciatus	15	NANA2	2	108.80				
C	Chaetodon lunula	15	NANA2	2	71.99				
C	Plectroglyphidodon johnstonianus	15	NANA2	1	1.72				
C	Labroides phthirophagus	15	NANA2	1	0.36				
C	Thalassoma duperrey	15	NANA2	2	54.78				
C	Thalassoma duperrey	15	NANA2	3	164.85				
C	Thalassoma duperrey	15	NANA2	5	485.26				
C	Stethojulis balteata	15	NANA2	1	72.39				
C	Halichoeres ornatissimus	15	NANA2	1	25.14				
C	Halichoeres ornatissimus	15	NANA2	1	16.45	23	1214.49	3.7	
CF	Chaetodon unimaculatus	15	NANA2	2	28.45				
CF	Chaetodon multicinctus	15	NANA2	2	26.06				
CF	Chaetodon multicinctus	15	NANA2	1	2.80				
CF	Cantherhines dumerili	15	NANA2	1	301.33	6	358.63	1.1	
H	Acanthurus triostegus	15	NANA2	4	400.06				
H	Acanthurus leucopareius	15	NANA2	6	165.09				
H	Acanthurus leucopareius	15	NANA2	60	7962.02				
H	Acanthurus nigrofusus	15	NANA2	2	108.50				
H	Acanthurus nigrofusus	15	NANA2	15	358.46				
H	Acanthurus nigrofusus	15	NANA2	9	67.72				
H	Acanthurus olivaceus	15	NANA2	24	13516.09				
H	Acanthurus dussumieri	15	NANA2	9	1504.95				
H	Ctenochaetus strigosus	15	NANA2	6	91.19				
H	Ctenochaetus strigosus	15	NANA2	6	161.06				
H	Ctenochaetus strigosus	15	NANA2	4	30.30				
H	Zembrasoma flavescens	15	NANA2	6	10.13				
H	Zembrasoma flavescens	15	NANA2	2	18.97	153	24394.54	74.9	
O	Stegastes fasciolatus	15	NANA2	4	59.05				
O	Melichthys niger	15	NANA2	2	496.06	6	555.12	1.7	
P	Chromis hanui	15	NANA2	1	0.75				
P	Naso hexacanthus	15	NANA2	9	1716.70				
P	Naso hexacanthus	15	NANA2	35	4309.13	45	6026.57	18.5	
	TOTAL	15	NANA2	233	32549.35	233	32549.35	99.9999999	
C	Aulostomus chinensis	16	PIPE	1	12.84				
C	Scomberoides laysan	16	PIPE	1	149.61				
C	Caranx melampygus	16	PIPE	4	771.41				
C	Lutjanus kasmira	16	PIPE	4	762.59				
C	Lutjanus kasmira	16	PIPE	77	4416.14				
C	Monotaxis grandoculis	16	PIPE	1	13.74				
C	Mulloides vanicolensis	16	PIPE	120	6062.18				
C	Mulloides vanicolensis	16	PIPE	115	17051.21				
C	Parupeneus multifasciatus	16	PIPE	5	272.00				
C	Parupeneus multifasciatus	16	PIPE	2	310.84				
C	Forcipiger flavissimus	16	PIPE	2	18.30				
C	Forcipiger flavissimus	16	PIPE	1	2.53				
C	Plectroglyphidodon johnstonianus	16	PIPE	7	12.05				
C	Paracirrhites arcatus	16	PIPE	14	228.88				
C	Cirrhitops fasciatus	16	PIPE	1	15.63				
C	Thalassoma duperrey	16	PIPE	19	520.42				
C	Thalassoma duperrey	16	PIPE	51	2802.39				
C	Thalassoma duperrey	16	PIPE	59	5726.07				
C	Thalassoma purpureum	16	PIPE	1	525.30				
C	Gomphosus varius	16	PIPE	5	310.15				
C	Gomphosus varius	16	PIPE	3	67.80				
C	Stethojulis balteata	16	PIPE	5	361.95				
C	Macropharyngodon geoffroy	16	PIPE	3	17.24				
C	Macropharyngodon geoffroy	16	PIPE	2	37.26				
C	Anampses chrysocephalus	16	PIPE	4	10.51				
C	Halichoeres ornatissimus	16	PIPE	3	49.35				
C	Halichoeres ornatissimus	16	PIPE	2	50.27				
C	Zanclus cornutus	16	PIPE	1	104.16				
C	Sufflamen bursa	16	PIPE	2	171.75	515	40854.57	69.0	
CF	Chaetodon multicinctus	16	PIPE	4	26.63				
CF	Chaetodon multicinctus	16	PIPE	2	26.06				
CF	Pervagor spilosoma	16	PIPE	2	32.43				
CF	Pervagor melanocephalus	16	PIPE	1	9.75	9	94.87	0.2	
H	Scarus sordidus	16	PIPE	4	3100.27				
H	Scarus sordidus	16	PIPE	5	700.03				
H	Scarus sordidus	16	PIPE	5	1835.11				
H	Scarus psittacus	16	PIPE	1	1437.46				
H	Acanthurus nigrofusus	16	PIPE	20	284.25				
H	Acanthurus nigrofusus	16	PIPE	15	358.46				
H	Ctenochaetus strigosus	16	PIPE	9	241.60	59	7957.18	13.4	
O	Stegastes fasciolatus	16	PIPE	16	415.68				
O	Melichthys vidua	16	PIPE	2	398.06				
O	Melichthys vidua	16	PIPE	2	139.84				
O	Canthigaster jactator	16	PIPE	7	24.93				
O	Canthigaster rivulata	16	PIPE	1	22.05	28	1000.57	1.7	
P	Chaetodon miliaris	16	PIPE	17	359.86				
P	Dascyllus albisella	16	PIPE	12	37.06				
P	Abudefduf abdominalis	16	PIPE	115	3603.25				
P	Abudefduf vaigensis	16	PIPE	124	3885.24				
P	Chromis ovalis	16	PIPE	125	1444.96	393	9330.36	15.8	
	TOTAL	16	PIPE	1004	59237.55	1004	59237.55	100	